

EVALUATION OF THE EFFECT OF PERCEPTION
OF RISK MESSAGES ON OBSERVED
SAFETY BELT USAGE

Patricia F. Waller
Livia K. Li
J. Richard Stewart
Juliana M. Ma

December 1983

~~HSRC-PR132~~

UNC/HSRC-83/12/2

Evaluation of the Effect of Perception of Risk Messages
on Observed Safety Belt Usage

Patricia F. Waller
Livia K. Li
J. Richard Stewart
Juliana M. Ma

The University of North Carolina
Highway Safety Research Center
Chapel Hill, NC 27514

Contract No. DTNH22-82-C-17271
Contract Amount: \$103,309

December, 1983

Final Report

Prepared for

U.S. DEPARTMENT OF TRANSPORTATION
NATIONAL HIGHWAY TRAFFIC SAFETY ADMINISTRATION
WASHINGTON, DC 20590

NOTICE

This document is disseminated under the sponsorship of the Department of Transportation in the interest of information exchange. The United States Government assumes no liability for the contents or use thereof.

1. Report No.	2. Government Accession No.	3. Recipient's Catalog No.	
4. Title and Subtitle Evaluation of the Effect of Perception of Risk Messages on Observed Safety Belt Usage		5. Report Date December, 1983	
		6. Performing Organization Code	
		8. Performing Organization Report No.	
7. Author(s) Patricia F. Waller, Livia K. Li, J. Richard Stewart, and Juliana M. Ma		10. Work Unit No. (TRAIS)	
9. Performing Organization Name and Address The University of North Carolina Highway Safety Research Center CPT 197A Chapel Hill, NC 27514		11. Contract or Grant No. DTNH22-82-C-17271	
		13. Type of Report and Period Covered Final Report October, 1982 - November, 1983	
12. Sponsoring Agency Name and Address Department of Transportation NHTSA Washington, DC		14. Sponsoring Agency Code	
15. Supplementary Notes			
16. Abstract <p>This study examined the effect of carefully developed safety belt messages on observed belt usage. Subjects were divided into six groups, including two control groups, with different groups receiving different messages and/or treatments. All subjects appeared for three different sessions. Questionnaires asking about attitudes and behaviors in relation to various health problems and practices were administered at all meetings, but the message was presented only at the second meeting. Between the second and third meetings, subjects made a number of telephone call-ins and heard their specific message repeated. One group experienced a delay between the end of the telephone call-ins and the final meeting. Observations were made of belt usage as subjects arrived at and left the meetings.</p> <p>The overall findings were disappointing in that there was no observable effect of the messages on observed safety belt usage. Furthermore, there were no significant differences in observed belt usage as a function of age, race, or sex. Observed belt usage was associated with greater reported frequency of eye examinations but was not related to other reported health maintenance behaviors. "Saturation" of the message via telephone calls to listen to the message did not lead to increases in belt usage.</p> <p>On the basis of this study it cannot be concluded that brief messages on the effects of safety belt usage and the risk of motor vehicle injury, combined with follow-up messages via telephone, lead to increases in observed belt usage.</p>			
17. Key Words Seat Belts Safety Belts Risk Perception		18. Distribution Statement	
19. Security Classif. (of this report)	20. Security Classif. (of this page)	21. No. of Pages	22. Price

ACKNOWLEDGEMENTS

A large number of people participated in the conduct of this project. Although it would not be feasible to mention all of them by name, the authors wish to acknowledge those who were especially helpful.

George Chi prepared the original and revised experimental designs and procedures.

George Taylor, manager of the NC Memorial Hospital parking deck, kindly assisted in arrangements for obtaining a parking booth, and Robert Daniel arranged for setting up the booth and other necessary equipment.

Janice Scroggs arranged for the use of the Homestead Community Center and took responsibility for seeing that it was adequately equipped for the purposes of the project.

Laurence Naumoff, William Stephenson, and Sam Sockwell assumed major responsibility for running subjects, assisted by Michael Gilmartin, Carol Popkin, and Robert Gray. Michael Gilmartin also assumed primary responsibility for data entry. Their conscientious and reliable performance was essential to the project, and they were all a joy to work with.

Atwood Askew and Julie Glasser of Belk Legget, a local department store, were most cooperative in providing a selection of incentive gifts at a bulk rate price to the project.

Linda Snyder of Piedmont Airlines arranged for the project to receive, gratis, a supply of packaged toasted almonds to distribute to subjects at each session. We are most grateful for such generosity.

Judy Hall, Peggy James, and Teresa Parks cheerfully provided secretarial throughout the project.

Bruce Bigelow served as contract manager during the first half of the project, when he was replaced by Michael Smith. Both have been patient and supportive throughout a difficult project.

Finally, the authors wish to express their gratitude to the many people who drove long distances and gave of their time to serve as subjects in this study.

To all of these people the authors express their appreciation.

EXECUTIVE SUMMARY

Although it is well known that increased safety belt usage would lead to a significant decrease in motor vehicle injuries and deaths, actual belt usage remains very low. Most government sponsored efforts still focus on voluntary rather than mandatory approaches. Part of this larger effort includes developing and testing public information materials.

This study examined the effect of carefully developed safety belt messages on observed belt usage. Subjects were divided into six groups. The first three each received one of three safety belt messages in audiovisual form. Group 4 received the same message as group 3 but in audio form only. Groups 5 and 6 received the same message, one on Heart and Exercise in audiovisual form. Groups 1 through 5 received an experimental questionnaire that asked about a variety of health-related attitudes and behaviors, including motor vehicle accidents and safety belt usage. Group 6 received a control questionnaire that substituted questions on blood pressure and stroke for the motor vehicle injury and safety belt questions, but all other questions were the same as those on the experimental questionnaire.

All subjects appeared for three different sessions. At the first session the questionnaire was administered. At the second session the message was presented, followed by the questionnaire. At the third session only the questionnaire was administered. Between the second and third sessions each subject made a total of 15 telephone calls to listen to a pre-recorded audio version of the message appropriate to his group.

All subjects came for their last session in the week following their last telephone call-in except for those in group 2. These subjects experienced a two-week delay between the week of their last telephone call and their final meeting.

Observations were made of safety belt usage as subjects arrived for the sessions and again as they left. For participation in the study, subjects were given an incentive, namely, a gift they could select from a variety of merchandise or \$25 in cash or a gift certificate from a local department store. In addition, for each meeting attended each subject was given a raffle ticket that made him eligible for prizes in a drawing conducted after the final meetings had been held. To be eligible for the grand prize, a subject had to have attended all meetings and complete all telephone call-ins. To be eligible for the lesser prizes, a subject still had to have completed all meetings and at least 12 of the 15 required telephone call-ins.

The overall findings were disappointing in that there was no observable effect of the messages on observed safety belt usage. In-depth analyses of the data were conducted to identify any possible relations or leads for further investigation.

There were some differences in the proportions of belt wearers according to age, sex, and race, but none of the differences was statistically significant.

When reported frequency of belt usage was related to reports of other health maintenance behaviors, there was a relationship to frequency of dental flossing, as well as a weaker relationship to frequency of eye examinations. Observed belt usage was also related to reported frequency of eye examinations, suggesting that the finding may be real. The relationship between reported belt usage and observed belt usage was extremely significant.

The group receiving the message in audio form only showed significantly less improvement than the group that received the same message in audiovisual form. However, the interpretation of this finding is problematical because the group receiving the particular message in audiovisual form was also significantly better than the other two message groups that received audiovisual presentations. It is possible that the particular message used (Physics) was much superior to the other two messages and that indeed the audiovisual presentation was more effect, but with the relatively small numbers of subjects in each group and the large number of statistical tests conducted, it may not be prudent to accept this interpretation without further investigation.

No differences were found between subjects experiencing a delay between their last call-in message and the final meeting and subjects whose last meeting was conducted immediately after the call-ins were completed. Thus there was no evidence of a delay either "consolidating" the impact of the message or contributing to a weakening of message effect. The additional contributions of the call-in messages were not related to any increment in observed belt usage. Thus it cannot be concluded that the additional exposure to the messages reinforced or strengthened the tendency to wear belts.

During the conduct of the study, a community safety belt program was instituted, with the initial launching occurring just three weeks prior to the first meeting sessions of this study and continuing throughout the duration of this project. Measures of awareness of the community safety belt project did not indicate that it influenced the basic findings of this study. However, the possibility cannot be ruled out that this other activity in some way affected the results of this project.

Subjects' reported judgments of the three major health problems in the U.S. today did not appear to be differentially affected by the various messages presented on safety belts, but the groups receiving safety belt messages showed a greater increase in the perception of accidents (but not necessarily motor vehicle) as a major health problem.

The basic findings are listed below:

1. There were no statistically significant differences in observed belt usage as a function of age, race, or sex.
2. Observed belt usage was associated with greater reported frequency of eye examinations but was not related to other reported health maintenance behaviors included in the questionnaire.

3. There were no significant relationships found between the safety belt messages and observed belt usage.

4. Although the comparison of the message effects of audiovisual presentation versus audio presentation were statistically significant, the finding was not considered valid because the particular audiovisual group in question showed significantly better results than the other audiovisual message groups in the study.

5. "Saturation" with the message via telephone calls to listen to the message did not lead to increases in belt usage.

6. There was no evidence that the Community Safety Belt Program affected the major findings of the study.

7. The groups receiving safety belt messages showed a greater increase in the proportion of subjects reporting accidents as a major health problem.

On the basis of this study it cannot be concluded that brief messages on the effects of safety belt usage and the risk of motor vehicle injury, combined with follow-up messages via telephone, lead to increases in observed belt usage.

TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENTS.....	iii
EXECUTIVE SUMMARY.....	v
LIST OF TABLES.....	xi
INTRODUCTION.....	1
METHOD.....	3
Experimental Design.....	3
Subjects.....	5
Recruitment.....	5
Scheduling Procedures.....	6
Incentives.....	6
Meeting Site.....	7
Questionnaires.....	8
Data Collection.....	9
The First Session.....	9
The Second Session.....	10
Telephone Call-Ins.....	10
The Third Session.....	11
Data Editing and Creation of Analysis File.....	12
RESULTS AND DISCUSSION.....	13
Overall Belt Usage Rates.....	13
Belt Usage Rates by Demographic Groups.....	13
Belt Usage by Race.....	13
Belt Usage by Sex.....	13
Belt Usage by Age.....	14
Belt Usage by Other Health Maintenance Behaviors.....	14
Effect of Message.....	15
Audiovisual Versus Audio Presentation.....	18
Effect of Delay.....	18
Effect of Saturation.....	19
Effect of Community Safety Belt Project.....	19
Perceived Health Problems.....	21
SUMMARY AND CONCLUSIONS.....	22
APPENDICES	
1. Description of Safety Messages.....	25
2. Original and Revised Experimental Designs.....	27
3. Information Form Completed for Potential Subjects.....	29
4. Schedule Information Sent to Subjects.....	31
5. List of Incentive Gifts.....	33
6. Parking Booth Arrangement.....	35

TABLES OF CONTENTS (Continued)

	Page
7. Health Attitude and Behavior Questionnaires.....	37
8. Subject Check-In Form.....	79
9. Information Forms for Telephone Call-Ins.....	81
10. Information Form Explaining Incentive Raffle.....	87
11. Tables of Preliminary Cross-Tabs of Questionnaire Responses, Subject Demographics, Changes in Belt Usage, and Message Groups.....	89
REFERENCES.....	147

LIST OF TABLES

	Page
Table 1. Summary of Subjects at Different Stages of the Study.....	6
Table 2. Belt Usage Rates by Observation by Group.....	15
Table 3. Belt Usage by Race.....	14
Table 4. Belt Usage by Sex.....	16
Table 5. Belt Usage by Age.....	16
Table 6. Belt Status by Group (Inbelt 3).....	17
Table 7. Belt Status by Group (Outbelt 3).....	20
Table 8. Belt Status by Awareness of CBSP (Inbelt 3).....	22
Table 9. Awareness of CSBP by Group.....	22
Table 10. Awareness of CSBP by Treatment Category.....	22
Table 11. Changes in Perception of Accidents as a Major Health Problem.....	23

Additional tables may be found in Appendix 11.

INTRODUCTION

There is overwhelming evidence that increased use of safety belts is by far the most cost effective measure available in highway safety today. Safety belts are installed in most seating positions in the vast majority of the vehicles on the highway today. Yet belt usage peaked at no more than 25 - 30 percent and has actually declined in recent years to a nationwide estimate of usage in the order of 12-15 percent. These conditions prevail in the face of evidence that the majority of people believe that seat belts save lives.

In recent years there has been a concerted effort to focus on the protection of infants and small children in car crashes. The groundswell of activity and concern that has been initiated in the United States probably represents the most significant public health movement to occur in this nation in decades. As of this writing, forty states and the District of Columbia have passed legislation relating to proper restraint of small children, and many more are introducing such legislation. While most state laws limit their occupant restraint requirements to infants and very young children, New York has enacted legislation that by 1987 will raise to ten the age below which occupant restraints are required. Business and industry are beginning to recognize their vested interests in encouraging protection of their employees and are implementing programs designed to increase belt usage. While the impact of these efforts has been far from complete, nevertheless the results must be considered strikingly successful for a voluntary approach. In one community wide effort observed usage rate has increased to over 40 percent (HSRC report, in preparation).

However, the fact remains that overall usage remains dismally low. Recent information from the United Kingdom (Ashton, Mackay, and Camm, 1983) shows that the enactment of a mandatory belt usage law increased usage from between 30 and 40 percent to 95 percent or higher. If we persist in the voluntary approach, much more must be known about how to motivate people to protect themselves from the risk of motor vehicle injury. One way to bring about such a change is to modify the perception of risk of motor vehicle injury.

This study was undertaken to investigate whether messages carefully designed to influence safety belt attitudes and behavior could bring about changes in reported safety belt attitudes and observed safety belt usage. The experimental messages used communicated messages that attempted to influence risk of perception of motor vehicle injury and provide information concerning effectiveness of safety belts in reducing risk. The study attempted to include a broad range of subjects rather than focus on college students. It also investigated the relationship between observed belt usage and subject age, race, and sex; mode of message presentation; reported beliefs and practices concerning other health problems and health-related behavior, as well as other variables described in more detail below.

The perception of risk messages used in this project were developed under a separate NHTSA contract conducted by Perceptronics. The

development of the messages is described in their reports (Slovic, Lichtenstein, MacGregor, Fischhoff, and Schwalm, 1983; Schwalm, and Slovic, 1983).

To accomplish the above objectives, subjects were recruited from the community and scheduled for three sessions at a specified project location. Between the second and third session subjects were required to make a total of 15 telephone calls to listen to a pre-recorded message. Different subject groups received different messages and in some cases different calling schedules. In all, there was a total of six different groups varying by message, mode of message presentation, and delay between final message and final session. The details of the procedures are described below.

METHOD

Experimental Design

The final experimental design used included six groups, four of which were exposed to the experimental message and two to the control message. In addition, each group received an experimental or control questionnaire. Appendix 1 describes the messages used. Specifically:

Group 1 received an audiovisual message on Relative Risk and the experimental questionnaire.

Group 2 received an audiovisual message on One-Third Probability Plus Alcohol Plus Control and the experimental questionnaire.

Group 3 received an audiovisual message on Physics of Crash Plus Alcohol Plus Control and the experimental questionnaire.

Group 4 received an audio message on Physics of Crash Plus Alcohol Plus Control and the experimental questionnaire.

Group 5 received an audiovisual message on Heart and Exercise and the experimental questionnaire.

Group 6 received an audiovisual message on Heart and Exercise and the control questionnaire.

The experimental questionnaire include questions about habits and attitudes concerning a number of health problems, including motor vehicle injuries and safety belt usage. The control questionnaire was identical except that it substituted questions concerning stroke for the safety belt questions.

Because of the small number of subjects responding to the various solicitation efforts, these six groups were compiled from the original 11 group design (Appendix 2). It was felt that the modified groupings would allow us to address the major questions with sufficient numbers (preferably about 40 subjects per group) in order to arrive at valid conclusions. Because of the reduction in number of groups, it was necessary to eliminate certain design issues, e.g., the efficacy of a parking attendant versus a van to collect belt usage data. Nevertheless, if the original 11 groups had been retained, there would not have been enough subjects in each group to address adequately any of the original questions.

Using the modified experimental design, the following hypotheses were addressed:

1. There is no relationship between the demographic characteristics of subjects and the relationships between safety belt usage and message presentation.

Belt usage rates of the different demographic groups were examined. In addition, changes in belt use were examined in relation to age, race, and sex.

2. There are no relationships between safety belt use and other health maintenance activities. Since both experimental and control questionnaires contained questions on other health-related matters such as eye examinations, dental hygiene, exercise, diet, and smoking, comparisons were made of belt usage attitudes and behaviors with responses to these other questionnaire items for all groups.
3. There is no message effect, that is, none of the three safety belt messages tested has an influence on subjects' belt wearing behavior or perceived risk of motor vehicle injury. This hypothesis was investigated by making comparisons among the four experimental groups and the two control groups, with special focus on those subjects who were not belted when they came in for their first meeting. In addition, comparisons were made between Groups 5 and 6 to determine whether questions concerning safety belts influence subsequent safety belt usage. Groups 5 and 6 received the identical message on Heart and Exercise, but differed on the questionnaire received.
4. The manner (audio or audiovisual) of message presentation has no effect on safety belt usage. The results of Group 3 were compared to those of Group 4 to address this hypothesis, since both groups received the same message (Physics) but in two different forms.
5. Observed belt usage is not modified as a function of time following last telephone message. The original 11-group design provided a better method of assessing the effect of delay than the modified design. Ideally, to measure the effect of delay unambiguously, there would be two identical message groups, one group attending the third meeting immediately after they finished their call-ins and the other group delaying their third meeting until two weeks after their last call-in. However, since this was not possible with the modified design, one message group, namely, Group 2, was dedicated to the delay question. Group 2 was compared to Groups 1, 3, and 4 on the basis of changes in belt usage rates between the outgoing belt usage at the second meeting and the outgoing belt usage at the third meeting. More, specifically, this analysis was restricted to those subjects who were not belted upon leaving the second meeting.
6. Saturation with the message does not produce any effect

above and beyond that obtained after the initial message presentation. Saturation was achieved by requiring subjects to call in to listen to a pre-recorded message five times a week for three weeks after the second meeting. Consequently, the effect of saturation for Groups 1, 3 and 4 was measured by comparing the outgoing belt usage of the second meeting with the incoming belt usage of the third meeting. For Group 2, that measure represented the effects of saturation and delay.

7. There is no influence on the study of the Community Safety Belt Project to encourage safety belt usage. Just prior to the first subject meeting in the present project, a major community wide program was launched to promote safety belt usage. The program included an extensive safety belt promotional campaign that used various type of incentives. To determine the effects of the community campaign on the results of this study, a question was included on each of the questionnaires asking about other health or safety messages heard in the previous week. This hypothesis was included to determine the number of subjects who knew about the community's safety belt incentive campaign so such comparisons could be made of belt usage by the knowledgeable subjects versus the naive subjects.
8. Participation in the study, including exposure to questions and messages concerning motor vehicle injuries, does not increase the perception of motor vehicle accidents as a health problem.

Subjects

Recruitment. Various efforts were undertaken to solicit subjects for participation in the study. Initially, subjects were solicited through their organizational affiliation, e.g., the North Carolina State Employees Association (through an announcement in their newsletter), local churches (both white and nonwhite), and the Retired Senior Volunteer Program (RSVP). These groups were chosen to provide a diverse population from which subjects could be recruited. However, insufficient responses were obtained from these organizations. As a result, additional recruitment efforts were undertaken. Flyers were distributed in town parking lots, and notices were posted around the Chapel Hill Community and in Hillsborough (in laundromats, banks, shops, courthouse, churches, around the University campus and other public places) to encourage participation. In addition, advertisements (including a large display ad purchased in a free semi-weekly newspaper) were placed in three local newspapers (The Daily Tarheel, the Chapel Hill Newspaper and the Village Advocate). Finally, special flyers were sent out to box holders of the three rural routes in the general vicinity of where the project was being conducted. With such extensive distribution and advertising efforts, 435 people responded by calling in for further information. However, almost 40 percent (172) of them dropped out prior to the first meeting. Reasons for dropping out were mostly because subjects were leaving town at the end of the school year (Table 1) or were unable to

Table 1. Summary of Subjects at the Different Stages of the Study.

Subjects responded to flyers and ads	435
No shows for the first meeting	172
No answer	40 (23.3%)
Left town	57 (33.1%)
Disconnected phone	26 (15.1%)
No Transportation	7 (4.1%)
Cancel by subject	36 (20.9%)
Miscellaneous	6 (3.5%)
No shows for the second meeting	8
Drop outs for the phone calls	2
No shows for the third meeting	5
Subjects who completed all phases of the study	248
Subjects who took the wrong questionnaire at the first meeting*	14
Subjects with records usable to the study**	234

* Removed from the analysis file

** Included those who came on bicycles, motorcycles, or walk-ins.

stay in town long enough to attend all three meetings. As a result, only 263 subjects agreed to participate in the study. Of these, 248 eventually completed.

Procedures for Scheduling Subjects. A special telephone line was set up to receive calls from subjects. The receptionist answered the call with "Health Research Study" and then obtained the following information from the caller: name, home telephone number, work telephone number, best hours to call, mailing address and how the subject heard about the project (Appendix 3). The caller was then told that a project staff member would call back soon to provide more information about the project and to set up appointments.

As calls came in, subjects were grouped according to how they had heard about the project. Within each such source group, they were randomly assigned to one of the original 11 treatment groups. However, the random assignment procedures were modified later because of differential drop out rates among the 11 groups. Thus, the later callers were assigned to groups by a weighted scheme, that is, groups with a higher dropout rate were assigned more people in order to keep the groups comparable in size. In addition, husbands and wives, family members, friends or co-workers were assigned to different groups. This was necessary to discourage outside discussion of the project during the duration of the study and to minimize carpooling. Carpooling tended to create a more demanding situation for obtaining belt usage information.

Once a subject was assigned to a group, efforts were made to contact the subject again to schedule the three meeting appointments. At the same time, subjects were informed of their necessary commitments to the study in return for a small appreciation gift. Ordinarily, meetings for the same group were scheduled during the same week, but subjects could usually pick a time within the week to accommodate their own schedules. Appointments were made on almost any day (even on legal holidays and on weekends, if necessary) from 7:00 in the morning to 6:00 in the evening. Following the phone call, a form confirming appointment dates and a map to the meeting site were sent to the subject (Appendix 4).

Efforts to contact subjects occurred continuously for several months of the study. At times, many calls were needed to locate a subject (40 subjects were "lost" because no contact could be made with them after numerous attempts). Additional calls were made to handle appointment changes and to remind subjects to make up their missed appointments or call-ins. In large part because of the continual monitoring of subjects' attendance, all but 15 of the subjects who came for the first meeting completed all phases of the study, representing almost a 95 percent completion rate for those that started in the study.

Incentives

Subjects were motivated to participate in the study by various methods. First, upon completion of the study, subjects were awarded with \$25 cash or gift certificate, or an item of their choice from a pre-selected list of merchandise. Arrangements were made with a local department store for obtaining a variety of merchandise for a discounted

price. All items (Appendix 5) offered by the store retailed for more than the bulk order price of \$25 (several retailed for as much as \$35). Samples of items were displayed during the first and second meetings, and subjects were requested to indicate their preference by the end of the second meeting. In this way, the desired number of each item could be obtained and the problem of having too few of any one selection could be avoided. Also, by offering a variety of items, there was something appropriate for almost every person in our diverse subject population. Upon completing the call-ins and the third meeting, a specially printed and sequentially numbered gift coupon was presented to the subjects for them to redeem the items of their choice at the participating department store.

In addition to the above, a two-tiered lottery system was set up to encourage subjects' continued participation. Each time they attended a meeting, they were given a raffle ticket that provided a chance to win extra gift certificates. The more meetings attended (up to three), the greater was the chance of winning something. As long as they completed 12 of the required call-ins, they could be included in the drawing for the gift certificates. Those that completed all three meetings and the entire 15 call-ins were included in the drawing for the grand prize, a microwave oven. The system was explained to subjects at the beginning of the study, so that they would know what they were required to do in order to be eligible to participate.

Twenty-four prizes were given in the drawing, including one microwave oven, three \$30 gift certificates, eight \$20 gift certificates and twelve \$10 gift certificates. Since there were 243 subjects completing the study, each subject had about one in ten chances of winning something in addition to the \$25 cash or gift. If a subject's name was drawn more than once, only the largest prize was awarded, so that no subject could receive more than one extra prize. The drawing of raffle tickets was conducted on July 15. Once the winners were notified, the list of winners was recorded on a telephone line and all subjects could call in to find out the results of the drawing. All subjects had previously been given the number to call for this information.

Considerable effort was made to insure that the subjects would find the sessions enjoyable. Soft music played in the background, and during the message sessions when it was necessary for some subjects to wait, magazines were provided for browsing. In addition, for all three sessions as subjects left the Community Center, each person in the car was given a package of toasted almonds, donated by Piedmont Airlines, and thanked for participating. This procedure also required that cars stop at the parking booth when exiting, thus facilitating the observation of belt usage.

Meeting Site

The Homestead Community Center was chosen as a meeting site because it met several important criteria. First, it is located about five miles out of town and has no local bus service so that it was highly likely that subjects would have to drive. Second, it is within reasonable commuting distance from the three surrounding communities: Chapel Hill, Hillsborough and Durham. Third, there is ample free parking space available. Fourth, it was possible to arrange to control entrance and egress. To accomplish

this, a sturdy parking booth of stainless steel construction, cones, and ropes (Appendix 6) were placed on the driveway outside the building to control entrance and exit of vehicles. Signs were installed on both sides of the drive to direct subjects into the Community Center's driveway. Fifth, there was adequate space in the building, including the possibility of having a waiting room, a testing room, and a message room. In addition, bathroom facilities were available.

The Community Center proved adequate on all these points. Most subjects arrived by car or pickup truck, although a few came by motorcycle or bicycle. The control of vehicles' entrance and exit enabled observation of belt usage in all vehicles. These observations were further enhanced by the "parking attendant's" signal for subjects to stop momentarily alongside the parking booth either for checking in (upon arriving) or for receiving a package of toasted almonds and turning in their raffle ticket (upon leaving).

Inside the building there were one large room, a kitchen, and a small entrance room. Drapes were installed to cover up the windows. Tables and chairs were set up in the large room where at times as many as seven or eight subjects at a time completed questionnaires. Additional space was used by children accompanying parents who were serving as subjects. Paper and colored pens were supplied to children to help keep them occupied. Smaller children and infants were held by project personnel while parents participated in the study. The kitchen was modified for presenting the messages in the second session. It was the darkest room in the building, and the audiovisual equipment could easily be set up on the counter.

Questionnaires

At each of the three meetings, subjects were asked to complete a 19-item questionnaire. The purpose of the questionnaires was to determine the subject's perception of accident/injury risk and to obtain information on his or her belt usage habits. Other questions asked about other types of health-related behaviors such as smoking, exercise, diet, dental habits, and eye examinations. Including these other questions enabled a comparison of belt usage behavior with other types of health maintenance behaviors (Appendix 7). Furthermore, it enabled the collection of belt usage information in a less obtrusive manner.

Two different types of questionnaire were used. Control questionnaires differed from the experimental questionnaires in the following manner: instead of having safety belt questions, they contained questions on stroke and blood pressure. The questionnaires for the three meetings (Forms 1, 2 and 3 for the Experimental Groups and Forms 4, 5 and 6 for the Control Group) were essentially identical except that for the second meeting there was an additional question (#10) that asked about the subjects' reactions to the message just presented. Having all other questions the same from the first through the third meetings provided measures of change (especially for the safety belt and risk perception questions) as a result of being presented with the messages.

Another question asked about health or safety related messages heard by the subject in the last week prior to completing the questionnaire.

This question was included in an attempt to detect the effect, if any, of an undertaking that coincided with this project. A community safety belt incentive program was initiated three weeks prior to the first session meetings in this project and continued throughout this project's duration.

The last question on the questionnaire asked for the subject's opinion as to the three most important health problems in the United States. The objective here was to determine how often subjects view motor vehicle accidents as an important health problem, and whether their opinion changed after viewing/hearing the messages.

The six different forms of the questionnaire were specially coded and produced in three different colors (one for each meeting) so that the proper questionnaire could be given to the right subjects at the appropriate time.

However, as will be noted later, even with such careful precautions, some subjects were administered an incorrect questionnaire.

Data Collection

The First Session. The first series of meetings were conducted during the second and third weeks of May. The purpose of the first session was to describe to subjects in detail the reason for the study (to evaluate different types of health and safety messages), to explain the requirements for their participation in return for a small gift and chances to participate in the lottery, and to obtain baseline measures of their risk perception and belt usage. They were then given a questionnaire to complete. Following completion of the questionnaire, subjects were asked to examine the display of gifts and be prepared to state their selections at the next meeting. If they were already decided, they could sign up immediately. Prior to their leaving, they were asked to fill in their names and addresses on the raffle ticket stub and turn it in to the parking attendant on their way out. They were also admonished to discuss the study with no one else until after the last session.

An unobtrusive method was used for observing and recording subjects' belt usage. As subjects drove up the driveway, the parking attendant stopped them to check them in and, at the same time, noted their belt usage. As subjects drove out, the parking attendant took their raffle ticket, handed them a package of toasted almonds, thanked them for coming and again noted their belt usage.

A special check-in list (Appendix 8) was used by the parking attendant for recording the incoming belt usage observations. Subjects' names were written on this list, together with the times they were scheduled to appear. Next to each subject's name were four boxes for coding the belt usage information. DB stands for Driver Belted, and DN stands for Driver Not Belted. The other two categories were reserved for passengers only: Passenger Belted (PB) and Passenger Not Belted (PN). The coding scheme used made no differentiation on passengers by their seat position. Although this information would have been helpful, obtaining it would have placed an extra burden on an already demanding observation and coding task.

Usually, appointments were scheduled on the hour with as many as seven

or eight appointments scheduled for the same time. The parking attendant was the only person outside to check people in and out and to observe and code belt usage. Outgoing belt usage was classified initially by placing the ticket stub in one of four compartments in a box inside the parking booth. The compartments were labelled DB, DN, PB and PN, corresponding to the categories described earlier. At the first convenient break, the appropriate label was placed on the back of each stub.

In addition to checking subjects in and out, the parking lot attendant made note of no-shows and communicated this information as soon as possible to the research personnel inside the building. As soon as possible an attempt was made to contact each no-show and reschedule the subject at the earliest convenience.

The Second Session. During the last two weeks of May the messages were presented to the subjects. As subjects arrived, the parking lot attendant gave each one a card that showed their name, group number, and color code. The attendant then directed the subjects to a front room where they were to wait until the project staff came for them. As soon as possible, a project staff member greeted them and brought them into the room (kitchen) where the audiovisual equipment was set up. By noting the group number, as well as the color code, on the subject's card, the project staff member then presented the appropriate message. Subjects in the same group arriving at the same time were presented with the message simultaneously. The message on Relative Risk was presented to Group 1. The message on One-Third Probability (plus Alcohol plus Control) was presented to Group 2. The message on Physics of Crash (plus Alcohol plus Control) was presented to Groups 3 and 4 with Group 4 receiving only the audio version. Finally, Groups 5 and 6 were presented the message on Heart and Exercise, obtained from the American Heart Association.

At the end of the message, subjects were given the questionnaire and the raffle ticket and asked to complete them before leaving. After completing the questionnaires, they were again asked to look at the gift display and note their choice on a tablet available for that purpose. Printed information regarding the required telephone calls was given to subjects at this time. These information sheets were color coded by Group Number and the telephone numbers given had been selected for easy recall. For example, the number for the groups receiving the Heart and Exercise message was 962-4278, or 962-HART (see Appendix 9). The procedures for collecting incoming and outgoing belt usage information described previously were used in this session as well.

Telephone Call-ins. Most of June was reserved for subjects making their telephone call-ins to hear the pre-recorded message. As a participation requirement, each subject was to call in and listen to the message five times a week for three weeks. The grand prize lottery was only for those who called in all 15 times to participate. Calls could be made any time of the day or night. However, each call had to be made on a separate day in order to count, that is, a subject could not make five calls in one day and have them count as five out of the 15 required calls. Since only five call-ins were required for each week, subjects could miss one or two call-ins during the week and still make it up during the weekends. Arrangements were also made to receive collect calls from out of

town to accomodate subjects on vacation or otherwise away. Subjects called from as far away as California to meet the study requirements.

The call-ins were continuously monitored by the project staff, and reminder phone calls were made if subjects fell behind schedule or called in on the wrong line. Thus, all kinds of efforts were undertaken to help subjects meet the project requirements. Furthermore, the promised gift was granted as long as subjects called in only 12 out of the required 15 times although these subjects were not eligible for the grand prize in the drawing. Because of the above efforts, only a small proportion (5%) of subjects failed to complete the study.

Special telephone equipment was purchased for handling the messages as well as the instructions that preceded and followed the message. Subjects called in through their designated telephone line (one telephone line for each message, with the telephone number given to subjects in a handout at the end of the second meeting), listened to the message, and then left their name and date of calling. All telephone lines were checked each day. Data tapes were removed from the telephone machines and the data were transcribed onto hard copy forms by two persons, each working independently from the other. This method of coding the telephone data provided a reliability check on the coding. Discrepancies identified were then checked against the telephone tapes again. Enough tapes were purchased so that all telephone data could be preserved until the end of project. In this way, questions arising about the number of times a subject had called in were easily checked against the original data on the tapes which contained a record of subjects' names and call-in dates.

The Third Session. Except for Group 2, subjects in all groups were asked to come in for their third meetings the week immediately following their last call-in. Subjects in Group 2 were scheduled to come in during the third week after their last call-in. Consequently, it took three weeks (from June 20 to July 8) to bring all subjects in for their third meetings.

At this last meeting, belt usage information was obtained using the procedures described above. When subjects first arrived, they were asked the number of times they had called in. This reported number was checked against the hard copy record. Of all those (255) that attended the second meeting, only two subjects did not complete the required 12 call-ins.

As before, subjects were asked to complete the questionnaires and the raffle tickets. Upon completion, they were given the gift of their choice. For those electing cash, a check of \$25 was given. Gift coupons were prepared ahead of time and given to those electing a merchandise gift or the store's gift certificate. These gift coupons were sequentially numbered and made tamper proof (printed in green ink over white paper) and each was signed personally by a project staff member. Receipts for the gift coupons or checks were obtained from all subjects. All these measures were taken so as to account for the distribution of gifts. Subject's name and the choice of gift were written on the face of each gift coupon. They were then instructed to take the gift coupons down to the participating department store to redeem the gift of their choice.

Upon departure, subjects were thanked for their participation and an

information hand out (Appendix 10) explaining the upcoming lottery was provided to them. Because there was still a need to observe their outgoing belt usage for this meeting, it was not possible to reveal the real purpose of the study to subjects at this time. Instead, subjects were told that they would be sent a short summary of the study after the study report is finished.

Data Editing and Creation of Analysis File

During the period when meetings were conducted, the belt usage information (from check-in list and raffle ticket) were added to the questionnaire data at the end of each day. Such a procedure was used so that merging of the questionnaire and behavior data could be accomplished with relative ease. Furthermore, it simplified the merging of the first, second, and third meeting data because it involved matching of three instead of nine sources of data. During the period of data collection, data were also being entered onto the Apple IIe computer for creation of an analysis file. A total of 248 sets of questionnaires were entered and matched, collating data for first, second, and third meetings. However, 14 subjects were administered an incorrect questionnaire, usually experimental subjects taking control questionnaires, at the first meeting. Thus, for these 14 subjects, there were no baseline measures on their risk perception and seat belt attitudes. For this reason, they were removed from all subsequent analysis leaving only 234 subjects' data in the analysis file. In addition, subjects took the wrong questionnaires (also with experimental subjects taking control questionnaires) during the second meeting. These subjects were retained in the analysis file but a special flag was put on their record. Their data could be used for most analyses. No special effort was made to remove them from the remaining analyses because the lack of significant findings elsewhere suggested it would not be worthwhile to re-analyze the data eliminating these subjects.

RESULTS AND DISCUSSION

Overall Belt Usage Rates

Baseline belt usage rates (Inbelt 1) vary widely among the six message groups, from a low of 30 percent for the Relative Risk Group to a high of 57 percent for the group receiving the message on Physics of Crash (Table 2). Except for Group 5, all message groups showed a steady increase in belt usage from the first through the third meetings. However, this increase probably could not be attributed to the safety belt message heard because a control group (6) also showed a similar amount of increase. Statistical tests comparing the above results are reported in a later section of the report, and more detailed information can be found in Appendix 11. However, because of small N's and large variances, many absolute differences were not statistically significant.

Belt Usage Rates by Demographic Groups

Belt Usage by Race. Higher belt usage rates were observed for the white drivers for all three meetings (Table 3). A higher proportion of nonwhite drivers was found to be using belts at the third meeting as compared to the first meeting, and this increase was comparable to that observed for the white drivers. However, because of the lower initial usage rates of the nonwhite drivers, this change represented a larger amount of proportional increase for them, that is, almost a 50 percent increase for the nonwhite drivers as compared to a 32 percent increase for the white drivers. Nevertheless, neither the original observations of belt usage nor the changes in usage showed significant differences by race.

Table 3. Belt Usage by Race

	WHITE	NONWHITE
Inbelt 1	42.5%	28.0%
Outbelt 1	43.3%	28.0%
Inbelt 2	50.0%	40.0%
Outbelt 2	52.8%	40.0%
Inbelt 3	55.5%	44.0%
Outbelt 3	56.0%	41.7%
TOTAL N (At Inbelt 1)	201	25

Belt Usage by Sex. Females were found to have a higher initial belt usage rates but by the time they left the third meeting, the belt usage rates were very similar for both male and female subjects (Table 4). Thus, male subjects showed a higher proportional increase in belt usage than their female counterparts. However, neither the original observations of

Table 2. Belt Usage Rates by Groups (Percent)

TRIP	GROUP					
	1 Relative Risk	2 1/3 Alcohol + Control	3 Physics,Alc., Control	4 Physics,Alc., Cont.,Audio	5 Heart, Belt ?	6 Heart, Control
Inbelt 1	30.0	52.6	57.5	33.3	40.0	31.8
Outbelt 1	27.5	57.9	60.0	36.4	40.0	29.5
Inbelt 2	38.5	55.3	65.0	42.4	57.1	39.5
Outbelt 2	41.0	65.8	65.0	51.5	54.3	34.9
Inbelt 3	48.7	62.2	69.2	45.5	48.5	50.0
Outbelt 3	46.2	64.9	69.2	54.5	42.4	47.6

belt usage nor the change in belt use were statistically significant by sex.

Table 4. Belt Usage by Sex

	MALE	FEMALE
Inbelt 1	35.2%	42.7%
Outbelt 1	35.2%	43.6%
Inbelt 2	47.2%	49.4%
Outbelt 2	43.4%	53.8%
Inbelt 3	47.2%	56.4%
Outbelt 3	54.7%	54.4%
TOTAL N (at Inbelt 1)	54	172

Belt Usage by Age. The belt usage data by age are quite consistent across the six observation points. Subjects in the middle category (26-35) showed the highest belt usage rates, followed by subjects who were age 36 or above (Table 5). Younger subjects were found to be using belts less often than the other two age groups as they showed up for the meetings. However in terms of net change (Inbelt 3 - Inbelt 1), younger and older drivers showed a greater amount of change than drivers in the middle age category. Nevertheless, none of the observed age differences between or within age groups was found to be statistically significant.

Table 5. Belt Usage by Age

	AGES 18-25	AGES 26-35	AGES 36 AND OVER
Inbelt 1	34.3%	49.4%	37.0%
Outbelt 1	31.3%	51.2%	39.7%
Inbelt 2	42.4%	55.8%	46.6%
Outbelt 2	43.9%	60.5%	47.2%
Inbelt 3	51.5%	55.8%	54.8%
Outbelt 3	53.0%	62.4%	46.6%
TOTAL N (At Inbelt 1)	67	85	73

Belt Usage by Other Health Maintenance Behaviors

The questionnaires used included questions about a variety of health maintenance behaviors. The actual observed safety belt usage upon arrival

at the first meeting was related to the responses to questions concerning other health-related activities. There were no significant relationships found between observed belt usage and reported frequency of physical checkups, smoking behavior, frequency of exercise, or frequency of dental examinations. There was a significant relationship ($p < .03$) between observed belt usage and reported frequency of dental flossing, with belt users more likely to report frequent flossing. There was a weaker relationship ($p < .10$) found between observed belt usage and reported frequency of eye examinations. The tables for these findings, as well as others, may be found in Appendix 11.

Subjects' responses to the questions about belt usage were compared with other reported health maintenance behaviors. The only significant relationship found was between reported belt use and reported frequency of eye examination ($p < .005$). In addition, observed belt usage was compared with reported belt usage, and the relationship was highly significant ($p < .0001$), with persons who were observed to be wearing a belt much more likely to report frequent usage. Tables for these findings are also in Appendix 11.

Effect of Message

The effect of message was investigated based on each subject's safety belt wearing status when driving in for the third meeting (Inbelt 3) as contrasted with his status when driving in for the first meeting (Inbelt 1). This comparison maximized the possibility of detecting any effect, since by the third meetings subjects had had exposure to the message from both the second meeting and the telephone call-ins. More specifically, these analyses were restricted to subjects who were not wearing safety belts when they appeared for their first meeting. The response variable was, thus, belt wearing status (yes or no) on the third inbound trip.

An examination of whether age, race, or sex was related to change in belt usage by treatment groups showed no significant results (see III in listing of crosstabs in Appendix 11). Moreover, the χ^2 statistic for testing association between the response variable (Inbelt 3) and Group was likewise nonsignificant. Nonetheless, some simple categorical data models were fit to the Group by response frequencies to further characterize the variation in belt wearing rates by Group. The Group by response frequencies are shown in the following table (restricted to subjects who were not belted at the time of the first observation, Inbelt 1).

Table 6. Belt Status by Group (Inbelt 3)

GROUP	B	N	TOTAL
1	9	18	27
2	7	11	18
3	9	8	17
4	4	18	22
5	4	16	20
6	8	21	29

From this table a vector of the proportion of belted subjects by Group was computed and is given by:

$$P' = (.333 \quad .389 \quad .529 \quad .182 \quad .200 \quad .276)$$

To this vector is fit a model of the form:

$$\hat{P} = X\hat{B},$$

where X is a design matrix, \hat{B} is a vector of model coefficients estimated by a generalized least squares procedure, and \hat{P} is an estimator of P . In order to test some of the hypotheses of interest, the design matrix

$$X = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 & 1 \end{bmatrix}$$

was chosen. In this model $\hat{P} = X\hat{B}$,

$$\text{Where } B' = (b_1 \quad b_2 \quad b_3 \quad b_4 \quad b_5 \quad b_6)$$

the first component b_1 represents a weighted average belt wearing rate for Groups 1 and 2, since these two variables showed virtually the same proportions. The other components b_2, b_3, b_4, b_5 represent special effects for Groups 3-6, respectively, which either add to or subtract from the "baseline" (Groups 1 and 2) rate. Estimates of these effects are:

$$\begin{aligned} \hat{b}_1 &= .355 \\ \hat{b}_2 &= .175 \\ \hat{b}_3 &= -.173 \\ \hat{b}_4 &= -.155 \\ \hat{b}_5 &= -.079 \end{aligned}$$

This model fits the data quite well (as would be expected since only one degree of freedom is left). One hypothesis of interest involved a comparison of the belt usage rates for Groups 3 and 4 (represented by b_2 and b_3).

This hypothesis was tested by testing

$$H : b_2 = b_3$$

The χ^2 test of this hypothesis resulted in

$$\chi^2_1 = 5.64 \quad p = .0175,$$

so the rates for these two groups differ significantly. Other hypotheses of interest involve testing whether the control groups (Groups 5 and 6, indicated by b_4 and b_5 in the model, since Groups 1 and 2 are combined) differ significantly from the experimental groups. From the model shown

above, the special effects b_4 , and b_5 did not differ significantly from zero, and moreover, the last three effects (for Groups 4-6) could simultaneously be omitted from the model without significantly increasing the χ^2 due to error.

To more specifically compare the experimental and control groups a second model was fit to the data with

$$X = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 0 & 0 \\ 1 & 0 & 0 \\ 1 & 0 & 0 \\ 1 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix}$$

This model compares the belt wearing rates for Groups 5 and 6 with a weighted average rate for Groups 1-4. Neither of these groups differs significantly from the experimental groups.

Finally a model with

$$X = \begin{bmatrix} 1 & 0 \\ 1 & 0 \\ 1 & 1 \\ 1 & 0 \\ 1 & 0 \\ 1 & 0 \end{bmatrix}$$

was fit to the data. This model contains a weighted average rate for Groups 1, 2, 4, 5 and 6 and a special effect for Group 3. This model fits the data very well

$$\chi^2_4 = 3.30 \quad p = .51$$

and the special effect for Group 3 is significant ($p=.037$). In fact, the χ^2 due to error is so small that there can be no other one degree of freedom effect included in the model that would be significant at a .05 level of significance.

Nonetheless, special effects for Group 5 and for Group 6 were added to the model so that the hypothesis of equal effects for these groups could be tested. This procedure was to test whether or not the fact that Group 5 subjects took the safety belt questionnaire made a significant difference in changing their safety belt usage rates. The difference between these two effects was not statistically significant ($p = .53$).

In summary, these models show that the belt wearing rates for the control groups (presented with Heart and Exercise message) do not differ significantly from the average rate for the experimental groups (presented with the safety belt messages). Group 3 (presented with an audiovisual message on Physics of Crash) does have a significantly higher rate than the overall average for the other groups, and in particular the rate for Group 3 is significantly greater than the rate for Group 4 who were presented

with the same message on Physics of Crash but in an audio form only. This last analysis is discussed further below.

Audiovisual Versus Audio Presentation.

Group 3 received the Physics Plus Alcohol Plus Control message in audiovisual form, while Group 4 received the same message but in audio form only. As indicated in the section on Message Effect, the comparison between these two groups indicated that the audiovisual group increased significantly more in safety belt usage than did the audio only group. However, it is may not be valid to conclude that the audiovisual presentation is significantly superior to the audio presentation alone, since Group 3 appeared aberrant in its behavior and was significantly better than either of the other message groups. While it may be the case that the Physics message is vastly superior to the other two messages, the apparent similarity of content among the three messages suggests that the difference observed may not be attributable to the differences in the form of message presentation. The form of message presentation (audio versus audiovisual) should be investigated further, using a Physics message in comparison to other messages before conclusions are drawn.

The Effect of Delay

In order to investigate the effect of delay, the subjects' belt wearing status on leaving from the third meeting (Outbelt 3) was compared with that on leaving from the second meeting (Outbelt 2). More specifically, the analysis was restricted to those subjects who were not belted on leaving the second meeting and the response variable was taken as the belt status when leaving the third meeting. Group by response frequencies are shown in the following table:

Table 7. Belt Status by Group (Outbelt 3)

GROUP	B	N	TOTAL
1	4	19	23
2	2	10	12
3	4	10	14
4	2	14	16
5	4	12	16
6	6	20	26

The vector of proportions (belt wearing rates) on Outbelt 3 from this table is

$$P' = (.174 \quad .167 \quad .286 \quad .125 \quad .250 \quad .231)$$

A model with the design matrix

$$X = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 1 \end{bmatrix}$$

was first fit to the data. This model contains special effects for Group 2, Group 5, and Group 6 so that each of these groups could be compared with the weighted average rate of Groups 1, 3, and 4. None of these effects was statistically significant. Thus, delaying the third meeting for two weeks does not appear to make a difference on the Outbelt 3 rates for Group 2.

Effect of Saturation.

One of the questions of interest was whether the additional messages from the telephone call-ins served to reinforce any tendency to use safety belts. In order to test this possibility, safety belt usage for outbelt 3 was compared with that for Outbelt 2, and Groups 1, 3, and 4 (Relative Risk, AV; Physics, AV; and Physics, A) were compared with the control groups 5 and 6. The model described in the immediate prior section examined the changes from Outbelt 2 to Outbelt 3 for Groups 1, 3, and 4 as compared with Group 5 and again with Group 6. There were no significant differences found. Therefore it may be concluded that the saturation achieved by repeatedly exposing the subjects to the message via the telephone call-ins showed no effect.

Effect of Community Safety Belt Project

To assess the effects of the community safety belt project (CSBP) on the belt wearing behavior of the subjects in this study, the subjects were asked at each meeting to list any health or safety related messages they had heard during the previous week. The responses to these questions were categorized according to the nature of the message. One category corresponded to the Community Seat Belt project.

Of the 133 subjects who were not wearing safety belts when arriving for the first meeting, 20 indicated (at least once over the course of the three meetings) having heard a message concerning the Community Seat Belt Project. The remaining 113 did not indicate hearing such a message. Table 8 shows the belt wearing behavior of subjects arriving for the third meeting, divided by those who reported hearing of the Community Seat Belt Project and those who did not.

Table 8. Belt Status by Awareness of CSBP (Inbelt 3)

Awareness Status	Belted	Not Belted	Total
Heard of CSBP	9 (45.0%)	11 (55.0%)	20
Not Heard of CSBP	32 (28.3%)	81 (71.7%)	113
$\chi^2_1 = 2.22$ $p = .137$			

Table 8 shows that the belt wearing rate was higher for those subjects who had heard of CSBP than for those who had not, but with such small numbers this difference was not statistically significant. Table 9 shows the breakdown by treatment group of those who had heard of the CSBP and those who had not.

Table 9. Awareness of CSBP by Group

Awareness Status	1	2	3	4	5	6
Heard of CSBP	5 (18.5%)	3 (16.7%)	3 (17.7%)	4 (18.2%)	2 (10.0%)	3 (10.3%)
Not Heard of CSBP	22 (81.5%)	15 (83.3%)	14 (82.3%)	18 (81.8%)	18 (90.0%)	26 (89.7%)
Total	27	18	17	22	20	29

Table 9 shows that the subjects who had heard of the CSBP were quite evenly distributed across the six groups although the frequencies were slightly higher for the experimental groups than for the control groups. Finally, Table 10 shows a cross tabulation of awareness status by experimental and control groups.

Table 10. Awareness of CSBP by Treatment Category

AWARENESS STATUS	EXPERIMENTAL	CONTROL
Heard of CSBP	15 (17.9%)	5 (10.2%)
Not Heard of CSBP	69 (82.1%)	44 (89.8%)
Total	84	49

A χ^2 test of association on Table 9 yielded $\chi^2_1 = 1.42$, $p > .20$

While those subjects who reported awareness of the Community Seat Belt Project had slightly higher (but not significantly different) safety belt wearing rates at the time they appeared for the third meeting, the small number of these subjects and their even distribution across the study groups indicate that the CSBP should not produce any noticeable distortion in the results of the present study.

Perceived Health Problems

The questionnaires at each meeting asked the subjects to list what they considered to be the three most important health problems in the U.S. today. It was of interest to evaluate any changes that might have occurred between the first and third meetings with respect to the perception of accidents as a major health problem. To this end an analysis was conducted of subjects who listed accidents as a major health problems at the third meeting, but had not listed accidents at the first meeting. It should be noted that any reference to accidents was included, not solely motor vehicle accidents. Table 11 shows the frequencies of those who did and did not include accidents as a major health problem at the third meeting, given that they had not listed accidents at the first meeting.

Table 11. Changes in Perception of Accidents
as a Major Health Problem.

Mentioned Accidents	Group					
	1	2	3	4	5	6
Yes	4	9	7	5	3	0
	(12.50)	(29.03)	(21.21)	(16.13)	(10.34)	(0.00)
No	28	22	26	26	26	41
	(87.50)	(70.97)	(78.79)	(83.87)	(89.66)	(100.00)

A model was fit to these data which included special effects for each of the control groups (5 and 6) as compared with the experimental groups. A comparison between Groups 5 and 6 showed that they did not differ significantly. On the other hand, the two control groups combined did differ significantly from the experimental groups ($\chi^2_1 = 26.4$, $p = .0000$), with the experimental groups showing a greater increase in the proportions of subjects viewing accidents as a major health problem.

SUMMARY AND CONCLUSIONS

This study examined the effect of safety belt messages on observed belt usage. Subjects were divided into six groups. The first three each received one of three safety belt messages in audiovisual form. Group 4 received the same message as group 3 but in audio form only. Groups 5 and 6 received the same message, one on Heart and Exercise in audiovisual form. Groups 1 through 5 received an experimental questionnaire that asked about a variety of health-related attitudes and behaviors, including motor vehicle accidents and safety belt usage. Group 6 received a control questionnaire that substituted questions on blood pressure and stroke for the motor vehicle injury and safety belt questions, but all other questions were the same as those on the experimental questionnaire.

All subjects appeared for three different sessions. At the first session the questionnaire was administered. At the second session the message was presented, followed by the questionnaire. At the third session only the questionnaire was administered. Between the second and third sessions each subject made a total of 15 telephone calls to listen to a pre-recorded audio version of the message appropriate to his group.

All subjects came for their last session in the week following their last telephone call-in except for those in group 2. These subjects experienced a two-week delay between the week of their last telephone call and their final meeting.

Observations were made of safety belt usage as subjects arrived for the sessions and again as they left. For participation in the study, subjects were given an incentive, namely, a gift they could select from a variety of merchandise or \$25 in cash or a gift certificate from a local department store.

The overall findings were disappointing in that there was no observable effect of the messages on observed safety belt usage. In-depth analyses of the data were conducted to identify any possible relations or leads for further investigation.

There were some differences in the proportions of belt wearers according to age, sex, and race, but none of the differences was statistically significant. This finding was somewhat surprising because previous studies have consistently reported a race difference with nonwhites less likely to wear belts. However, there were few nonwhites in the study, despite extensive efforts to attract them, and those who participated tended to be in higher level occupations. Hence, nonwhite participants were probably even more atypical than were their white counterparts.

When reported frequency of belt usage was related to reports of other health maintenance behaviors, there was a relationship to frequency of dental flossing, as well as a weaker relationship to frequency of eye examinations. Observed belt usage was also related to reported frequency of eye examinations, suggesting that the finding may be real. The

relationship between reported belt usage and observed belt usage was extremely significant.

The group receiving the message in audio form only showed significantly less improvement than the group that received the same message in audiovisual form. However, the finding probably cannot be interpreted to mean that the audiovisual presentation is significantly better, because the group receiving that particular message in audiovisual form was also significantly better than any of the other message groups. It is therefore difficult to conclude that the difference observed was attributable to the form of message presentation.

No differences were found between subjects experiencing a delay between their last call-in message and the final meeting and subjects whose last meeting was conducted immediately after the call-ins were completed. Thus there was no evidence of a delay either "consolidating" the impact of the message or contributing to a weakening of message effect. The additional contributions of the call-in messages were not related to any increment in observed belt usage. Thus it cannot be concluded that the additional exposure to the messages reinforced or strengthened the tendency to wear belts.

During the conduct of the study, a community safety belt program was instituted, with the initial launching occurring just three weeks prior to the first meeting sessions of this study and continuing throughout the duration of this project. Measures of awareness of the community safety belt project did not indicate that it influenced the basic findings of this study. However, the possibility cannot be ruled out that this other activity in some way affected the results of this project.

Subjects' reported judgments of the three major health problems in the U.S. today did not appear to be differentially affected by the various messages presented on safety belts, but the groups receiving safety belt messages showed a greater increase in the perception of accidents (but not necessarily motor vehicle) as a major health problem.

The basic findings are listed below:

1. There were no statistically significant differences in observed belt usage as a function of age, race, or sex.
2. Observed belt usage was associated with greater reported frequency of eye examinations but was not related to other reported health maintenance behaviors included in the questionnaire.
3. There were no significant relationships found between the safety belt messages and observed belt usage.
4. Although the comparison of the message effects of audiovisual presentation versus audio presentation were statistically significant, the finding was not considered conclusive because the particular audiovisual group in question showed significantly better results than the other audiovisual message groups in the study. With the relatively small numbers of subjects in each group and the very large number of statistical tests

that were calculated in this investigation, it would be premature to conclude that the differences observed between the audio and audiovisual presentations are valid.

5. "Saturation" with the message via telephone calls to listen to the message did not lead to increases in belt usage.

6. There was no evidence that the Community Safety Belt Program affected the major findings of the study.

7. The groups receiving safety belt messages showed a greater increase in the proportion of subjects reporting accidents as a major health problem.

On the basis of this study it cannot be concluded that brief messages on the effects of safety belt usage and the risk of motor vehicle injury, combined with follow-up messages via telephone, lead to increases in observed belt usage.

Appendix 1. Description of Safety Messages

1. Relative Risk

You think Road Runner cartoons are silly? Here's something even more silly. You probably lock the door to your home without a second thought. Why? It's simple protection! Then why don't you buckle up for the same protection. It could save you from even worse problems, like getting badly hurt. So why do you lock your door and not wear a seat belt? And you think cartoons are silly! Buckle up and play it safe.

2. One in Three Probability

You think the Road Runner is hard on Wiley Coyote? Well, real life is not any easier on you. At some time during the 50,000 car trips you'll take in your lifetime, one out of every three of you who drive will suffer from a serious, possibly fatal, accident. And no matter how well you drive, you can't control the drunks and bad drivers on the road. But you are in control when you wear your seat belt. Buckle up and beat the odds!

3. Physics of Crash

You think Wiley Coyote gets into too many crashes. Maybe you do, too. In a car accident there's a crash when your car hits something and a second, even worse, crash inside the car when you hit the windshield. Now, you can't always control the first crash, especially with the drunks and bad drivers on the road. But you can stop the second crash. When you wear your seat belt, it just never happens. Buckle up and take control!

4. Heart and Exercise

I'm Fred Brown. Some folks call me Downtown because I take such long shots. That's on the court. Off the court, no one should take a long shot with his health. One way I exercise to keep my heart healthy is by jumping rope. The American Heart Association promotes rope jumping nationwide for old guys like me and for kids. It's a great do anywhere, anytime kind of exercise. Call your American Heart Association if you want to jump rope for your heart.

One of America's leading killers is heart and blood vessel diseases.

Revised Basic Experimental Design
(Original Design)

Approaches		Risk Perception		Conventional		Baseline	
Target Population	Threshold of Risk Tolerance	Certainty/Pseudocertainty	Relative Risk			Pre-intervention Questionnaire Non-seat Belt Related Message	Irrelevant Questionnaire Non-seat Belt Related Message
Group	M	M	M	M	M		
1	*(40)#						
2		*(40)					
3			*(40)				
4				*(40)			
5	†(40)						
6						*(40)	
7							*(40)
8		†(40)					
9						** (40)	
10			†(40)				
11				†(40)			

* = Audio-visual presentation + multiple audio presentations

† = Multiple audio presentation only

belt usage behavior observed & recorded by parking lot attendant.

** = Same as group 6, except belt usage behavior recorded by observers in van

= Sample size

Modified Experimental Design

Approaches	Experimental Groups			Control Groups	
	Relative Risk	One in Three	Physics	Heart, Exp, Quest.	Heart Cont' Quest.
Group					
1 & 8	*(40)#				
2 & 5		*(40)			
3			*(40)		
10			†(40)		
4 & 6				*(40)	
7, 9, 11					*(40)

* = Audio-visual presentation + multiple audio presentations

† = Multiple audio presentations only

= Sample size

HEALTH RESEARCH STUDY FORM

How did you hear about this project? _____

Thank you for calling.

Appendix 4. Schedule Information Sent to Subjects

Schedule for Health Research Study

Name: _____

1st Meeting _____


2nd Meeting _____

3rd Meeting _____

Between meeting 2 and 3 you will need to call in (call collect, if you are out of town) 5 times a week for 3 weeks. We will give you further instructions at the end of your second meeting.

You will receive your gift only after the entire schedule has been met.

Please call 962-6578 if you have any questions or problems meeting this schedule or if you need to change a meeting time.



Appendix 5. List of Incentive Gifts

Gift Selection

	<u>Retail</u>
Waring 14 sp. Futura II blender	\$29.99
Waring ice cream freezer	32.99
Ralph Lauren striped knit shirt	35.00
14K 16" gold chain	35.00
(74625) fm/am electronic digital clock radio	29.95
J.G. Hook button down shirt (men)	31.00
Fieldcrest Accent lace full size blanket	29.00
(66217) men's Timex watch	29.95
(18411) women's Timex Cavatina	28.95
Opium Perfume (1.2 oz.)	30.00
Nike "Intrepid" running shoe (men)	29.00
Shaeffer pen & pencil set (matte (black)	30.00
Hokey carpet sweeper	29.95
Monogrammed towel set (2 bath, 2 hand, 2 wash)	29.50
Nylon suit or dress bag	31.00
(R644/72) Hazel Attache	29.50
Gift Certificates	25.00

Appendix 6. Parking Booth Arrangement


Procedures for Collecting Safety Belt Usage

Safety belt usage data will be collected when subjects arrive at and leave the sessions. The attached diagram shows the arrangement for vehicles arriving and leaving.

An observer will be stationed in the parking booth and all vehicles will be required to pass the parking booth in order to enter the premises. Small stakes connected by fluorescent tape will prevent vehicles from entering any other way, and a large sign will be posted by the booth saying, "Check in Here." Subjects will give their name as they enter and check in. When this is done the observer will note whether belts are used and indicate this on the check-in list. Instructions will also be given as to where to park.

Just prior to subjects' leaving, they will be given a lottery ticket on which they are to fill in their name, address, and phone number. This ticket is to be given to the attendant at the parking booth when the subject leaves the premises. As the subject passes the parking booth, the attendant will stop the car, give the subject a packet of toasted almonds, and collect the lottery ticket. The ticket will be put into one of two boxes, depending on whether belts are used or not. This part of the data collection may become difficult if there are several subjects in the car and if some are belted while others are not. However, we will have to see how this works out.

Information on belt usage can later be recorded from the lottery tickets, complete with names, according to the box in which they were placed.



HEALTH STUDY QUESTIONNAIRE - 1

FOR RESEARCH ONLY - ALL YOUR ANSWERS WILL BE COMPLETELY CONFIDENTIAL

Please mark only one answer to each of the following questions:

1. How concerned are you about dying from a heart attack?

☐ 1 Not concerned☐ 2 Only a little concerned☐ 3 Somewhat concerned☐ 4 Quite a bit concerned☐ 5 Greatly concerned

2. How often do you visit your doctor for a physical checkup?

☐ 1 Less than once in three years☐ 4 About once a year☐ 2 About once in three years☐ 5 More than once a year☐ 3 About once in two years

3. How often do you use dental floss on your teeth?

☐ 1 Never☐ 2 A few times a month☐ 3 Two or three times a week☐ 4 Once a day☐ 5 More than once a day

4. How concerned are you about being crippled by an automobile accident?

1 Not at all concerned

2 Only a little concerned

3 Somewhat concerned

4 Quite a bit concerned

5 Greatly concerned

5. How often do you wear seat belts while driving?

1 Never

2 Rarely

3 Sometimes

4 Most of the time

5 Always

6. About how many cigarettes do you smoke each day?

1 None

2 Less than 1/2 a pack

3 1/2 to 1 pack

4 1 to 2 packs

5 More than 2 packs

7. Having lung cancer is just a matter of fate so smoking doesn't make that big a difference. How much do you agree or disagree with this statement?

1 Strongly agree

2 Slightly agree

3 Do not agree or disagree

4 Slightly disagree

5 Strongly disagree

8. How concerned are you about being crippled by a heart attack?

1 Not concerned

2 Only a little concerned

3 Somewhat concerned

4 Quite a bit concerned

5 Greatly concerned

9. About how often do you exercise?

1 Not at all or hardly ever

2 Several times a year

3 One or two times a month

4 Two or three times a week

5 Daily or almost daily

10. Having a heart attack is just a matter of fate, so proper diet and exercise don't make that big a difference. How much do you agree or disagree with this statement?

1 Strongly agree

2 Slightly agree

3 Do not agree or disagree

4 Slightly disagree

5 Strongly disagree

11. Getting killed or injured in a car accident is just a matter of fate, so seat belts don't make that big a difference. How much do you agree or disagree with this statement?

1 Strongly agree

2 Slightly agree

3 Do not agree or disagree

4 Slightly disagree

5 Strongly disagree

12. How often do you go to have your eyes checked?

1 Less than once in three years

2 About once in three years

3 About once in two years

4 About once a year

5 I do not go unless my eyes give me trouble

13. The chances of getting into an accident are so small that seat belts aren't really worth the trouble. How much do you agree or disagree with this statement?

1 Strongly agree

2 Slightly agree

3 Do not agree or disagree

4 Slightly disagree

5 Strongly disagree

14. How concerned are you about eating foods that will help keep you healthy?

1 Not at all concerned

2 Not very concerned

3 Not concerned one way or another

4 Somewhat concerned

5 Very concerned

15. How often do you go to a dentist to get your teeth checked?

1 Less than once in two years

2 About once in two years

3 About once a year

4 About once every six months

5 I do not go to the dentist except when my teeth bother me.

16. How concerned are you about being killed in an automobile accident?

1 Not at all concerned

2 Only a little concerned

3 Somewhat concerned

4 Quite a bit concerned

5 Greatly concerned

17. How concerned are you about getting lung cancer?

1 Not at all concerned

2 Only a little concerned

3 Somewhat concerned

4 Quite a bit concerned

5 Greatly concerned

18. Can you remember any health or safety messages you have heard during the last week?

1 No

2 Yes Please describe the type of message heard: _____

19. In your opinion, what do you think are the three most important health problems in the U.S. today?

1. _____

2. _____

3. _____

What is your name? _____

What is your address? _____

What is your home telephone number? _____

How old are you? _____

What is your sex? Male _____ Female _____

What is your race? White _____ Black _____ Other _____

How far did you go in school?

- _____ Less than high school
- _____ High school
- _____ More than high school

HEALTH STUDY QUESTIONNAIRE - 2

FOR RESEARCH ONLY - ALL YOUR ANSWERS WILL BE COMPLETELY CONFIDENTIAL

Please mark only one answer to each of the following questions:

1. How concerned are you about dying from a heart attack?

☐ Not concerned☐ Only a little concerned☐ Somewhat concerned☐ Quite a bit concerned☐ Greatly concerned

2. How often do you visit your doctor for a physical checkup?

☐ Less than once in three years☐ About once a year☐ About once in three years☐ More than once a year☐ About once in two years

3. How often do you use dental floss on your teeth?

☐ Never☐ A few times a month☐ Two or three times a week☐ Once a day☐ More than once a day

4. How concerned are you about being crippled by an automobile accident?

1 Not at all concerned

2 Only a little concerned

3 Somewhat concerned

4 Quite a bit concerned

5 Greatly concerned

5. How often do you wear seat belts while driving?

1 Never

2 Rarely

3 Sometimes

4 Most of the time

5 Always

6. About how many cigarettes do you smoke each day?

1 None

2 Less than 1/2 a pack

3 1/2 to 1 pack

4 1 to 2 packs

5 More than 2 packs

7. Having lung cancer is just a matter of fate so smoking doesn't make that big a difference. How much do you agree or disagree with this statement?

1 Strongly agree

2 Slightly agree

3 Do not agree or disagree

4 Slightly disagree

5 Strongly disagree

8. How concerned are you about being crippled by a heart attack?

1 Not concerned

2 Only a little concerned

3 Somewhat concerned

4 Quite a bit concerned

5 Greatly concerned

9. About how often do you exercise?

1 Not at all or hardly ever

2 Several times a year

3 One or two times a month

4 Two or three times a week

5 Daily or almost daily

10. How do you feel about the message you heard today? How much do you agree or disagree with the message?

1 Strongly agree

2 Slightly agree

3 Do not agree or disagree

4 Slightly disagree

5 Strongly disagree

11. Having a heart attack is just a matter of fate, so proper diet and exercise don't make that big a difference. How much do you agree or disagree with this statement?

1 Strongly agree

2 Slightly agree

3 Do not agree or disagree

4 Slightly disagree

5 Strongly disagree

12. Getting killed or injured in a car accident is just a matter of fate, so seat belts don't make that big a difference. How much do you agree or disagree with this statement?

1 Strongly agree

2 Slightly agree

3 Do not agree or disagree

4 Slightly disagree

5 Strongly disagree

13. How often do you go to have your eyes checked?

1 Less than once in three years

2 About once in three years

3 About once in two years

4 About once a year

5 I do not go unless my eyes give me trouble

14. The chances of getting into an accident are so small that seat belts aren't really worth the trouble. How much do you agree or disagree with this statement?

1 Strongly agree

2 Slightly agree

3 Do not agree or disagree

4 Slightly disagree

5 Strongly disagree

15. How concerned are you about eating foods that will help keep you healthy?

1 Not at all concerned

2 Not very concerned

3 Not concerned one way or another

4 Somewhat concerned

5 Very concerned

16. How often do you go to a dentist to get your teeth checked?

1 Less than once in two years

2 About once in two years

3 About once a year

4 About once every six months

5 I do not go to the dentist except when my teeth bother me.

17. How concerned are you about being killed in an automobile accident?

1 Not at all concerned

2 Only a little concerned

3 Somewhat concerned

4 Quite a bit concerned

5 Greatly concerned

18. How concerned are you about getting lung cancer?

1 Not at all concerned

2 Only a little concerned

3 Somewhat concerned

4 Quite a bit concerned

5 Greatly concerned

19. Can you remember any other health or safety messages you have heard during the last week?

1 No

2 Yes Please describe the type of message heard: _____

20. In your opinion, what do you think are the three most important health problems in the U.S. today?

1. _____

2. _____

3. _____

What is your name? _____

What is your address? _____

What is your home telephone number? _____

How old are you? _____

What is your sex? Male _____ Female _____

What is your race? White _____ Black _____ Other _____

How far did you go in school?

_____ Less than high school
_____ High school
_____ More than high school

HEALTH STUDY QUESTIONNAIRE - 3

FOR RESEARCH ONLY - ALL YOUR ANSWERS WILL BE COMPLETELY CONFIDENTIAL

Please mark only one answer to each of the following questions:

1. How concerned are you about dying from a heart attack?

☐ Not concerned☐ Only a little concerned☐ Somewhat concerned☐ Quite a bit concerned☐ Greatly concerned

2. How often do you visit your doctor for a physical checkup?

☐ Less than once in three years☐ About once a year☐ About once in three years☐ More than once a year☐ About once in two years

3. How often do you use dental floss on your teeth?

☐ Never☐ A few times a month☐ Two or three times a week☐ Once a day☐ More than once a day

4. How concerned are you about being crippled by an automobile accident?

1 Not at all concerned

2 Only a little concerned

3 Somewhat concerned

4 Quite a bit concerned

5 Greatly concerned

5. How often do you wear seat belts while driving?

1 Never

2 Rarely

3 Sometimes

4 Most of the time

5 Always

6. About how many cigarettes do you smoke each day?

1 None

2 Less than 1/2 a pack

3 1/2 to 1 pack

4 1 to 2 packs

5 More than 2 packs

7. Having lung cancer is just a matter of fate so smoking doesn't make that big a difference. How much do you agree or disagree with this statement?

1 Strongly agree

2 Slightly agree

3 Do not agree or disagree

4 Slightly disagree

5 Strongly disagree

8. How concerned are you about being crippled by a heart attack?

1 Not concerned

2 Only a little concerned

3 Somewhat concerned

4 Quite a bit concerned

5 Greatly concerned

9. About how often do you exercise?

1 Not at all or hardly ever

2 Several times a year

3 One or two times a month

4 Two or three times a week

5 Daily or almost daily

10. Having a heart attack is just a matter of fate, so proper diet and exercise don't make that big a difference. How much do you agree or disagree with this statement?

1 Strongly agree

2 Slightly agree

3 Do not agree or disagree

4 Slightly disagree

5 Strongly disagree

11. Getting killed or injured in a car accident is just a matter of fate, so seat belts don't make that big a difference. How much do you agree or disagree with this statement?

1 Strongly agree

2 Slightly agree

3 Do not agree or disagree

4 Slightly disagree

5 Strongly disagree

12. How often do you go to have your eyes checked?

1 Less than once in three years

2 About once in three years

3 About once in two years

4 About once a year

5 I do not go unless my eyes give me trouble

13. The chances of getting into an accident are so small that seat belts aren't really worth the trouble. How much do you agree or disagree with this statement?

1 Strongly agree

2 Slightly agree

3 Do not agree or disagree

4 Slightly disagree

5 Strongly disagree

14. How concerned are you about eating foods that will help keep you healthy?

1 Not at all concerned

2 Not very concerned

3 Not concerned one way or another

4 Somewhat concerned

5 Very concerned

15. How often do you go to a dentist to get your teeth checked?

1 Less than once in two years

2 About once in two years

3 About once a year

4 About once every six months

5 I do not go to the dentist except when my teeth bother me.

16. How concerned are you about being killed in an automobile accident?

- 1 Not at all concerned
- 2 Only a little concerned
- 3 Somewhat concerned
- 4 Quite a bit concerned
- 5 Greatly concerned

17. How concerned are you about getting lung cancer?

- 1 Not at all concerned
- 2 Only a little concerned
- 3 Somewhat concerned
- 4 Quite a bit concerned
- 5 Greatly concerned

18. Can you remember any health or safety messages you have heard during the last week?

- 1 No
- 2 Yes Please describe the type of message heard: _____
- _____
- _____

19. In your opinion, what do you think are the three most important health problems in the U.S. today?

1. _____
2. _____
3. _____

What is your name? _____

What is your address? _____

What is your home telephone number? _____

How old are you? _____

What is your sex? Male _____ Female _____

What is your race? White _____ Black _____ Other _____

How far did you go in school?

- _____ Less than high school
- _____ High school
- _____ More than high school

HEALTH STUDY QUESTIONNAIRE - 4

FOR RESEARCH ONLY - ALL YOUR ANSWERS WILL BE COMPLETELY CONFIDENTIAL

Please mark only one answer to each of the following questions:

1. How concerned are you about dying from a heart attack?

☐ Not concerned☐ Only a little concerned☐ Somewhat concerned☐ Quite a bit concerned☐ Greatly concerned

2. How often do you visit your doctor for a physical checkup?

☐ Less than once in three years☐ About once a year☐ About once in three years☐ More than once a year☐ About once in two years

3. How often do you use dental floss on your teeth?

☐ Never☐ A few times a month☐ Two or three times a week☐ Once a day☐ More than once a day

4. How concerned are you about being crippled by a stroke?

1 Not at all concerned

2 Only a little concerned

3 Somewhat concerned

4 Quite a bit concerned

5 Greatly concerned

5. How often do you get your blood pressure checked?

1 Never or rarely

2 Once every 4 or 5 years

3 Once every 2 or 3 years

4 Once a year

5 More than once a year

6. About how many cigarettes do you smoke each day?

1 None

2 Less than 1/2 a pack

3 1/2 to 1 pack

4 1 to 2 packs

5 More than 2 packs

7. Having lung cancer is just a matter of fate so smoking doesn't make that big a difference. How much do you agree or disagree with this statement?

1 Strongly agree

2 Slightly agree

3 Do not agree or disagree

4 Slightly disagree

5 Strongly disagree

8. How concerned are you about being crippled by a heart attack?

1 Not concerned

2 Only a little concerned

3 Somewhat concerned

4 Quite a bit concerned

5 Greatly concerned

9. About how often do you exercise?

1 Not at all or hardly ever

2 Several times a year

3 One or two times a month

4 Two or three times a week

5 Daily or almost daily

10. Having a heart attack is just a matter of fate, so proper diet and exercise don't make that big a difference. How much do you agree or disagree with this statement?

1 Strongly agree

2 Slightly agree

3 Do not agree or disagree

4 Slightly disagree

5 Strongly disagree

11. Having a stroke is just a matter of fate so keeping my blood pressure down doesn't make that big a difference. How much do you agree or disagree with this statement?

1 Strongly agree

2 Slightly agree

3 Do not agree or disagree

4 Slightly disagree

5 Strongly disagree

12. How often do you go to have your eyes checked?

1 Less than once in three years

2 About once in three years

3 About once in two years

4 About once a year

5 I do not go unless my eyes give me trouble

13. The chances of having a stroke are so small that getting my blood pressure checked really isn't worth the trouble. How much do you agree or disagree with this statement?

1 Strongly agree

2 Slightly agree

3 Do not agree or disagree

4 Slightly disagree

5 Strongly disagree

14. How concerned are you about eating foods that will help keep you healthy?

1 Not at all concerned

2 Not very concerned

3 Not concerned one way or another

4 Somewhat concerned

5 Very concerned

15. How often do you go to a dentist to get your teeth checked?

1 Less than once in two years

2 About once in two years

3 About once a year

4 About once every six months

5 I do not go to the dentist except when my teeth bother me.

16. How concerned are you about being killed by a stroke?

1 Not at all concerned

2 Only a little concerned

3 Somewhat concerned

4 Quite a bit concerned

5 Greatly concerned

17. How concerned are you about getting lung cancer?

1 Not at all concerned

2 Only a little concerned

3 Somewhat concerned

4 Quite a bit concerned

5 Greatly concerned

18. Can you remember any health or safety messages you have heard during the last week?

1 No

2 Yes Please describe the type of message heard: _____

19. In your opinion, what do you think are the three most important health problems in the U.S. today?

1. _____

2. _____

3. _____

What is your name? _____

What is your address? _____

What is your home telephone number? _____

How old are you? _____

What is your sex? Male _____ Female _____

What is your race? White _____ Black _____ Other _____

How far did you go in school?

- _____ Less than high school
- _____ High school
- _____ More than high school

HEALTH STUDY QUESTIONNAIRE - 5

FOR RESEARCH ONLY - ALL YOUR ANSWERS WILL BE COMPLETELY CONFIDENTIAL

Please mark only one answer to each of the following questions:

1. How concerned are you about dying from a heart attack?

☐ Not concerned☐ Only a little concerned☐ Somewhat concerned☐ Quite a bit concerned☐ Greatly concerned

2. How often do you visit your doctor for a physical checkup?

☐ Less than once in three years☐ About once a year☐ About once in three years☐ More than once a year☐ About once in two years

3. How often do you use dental floss on your teeth?

☐ Never☐ A few times a month☐ Two or three times a week☐ Once a day☐ More than once a day

4. How concerned are you about being crippled by a stroke?

1 Not at all concerned

2 Only a little concerned

3 Somewhat concerned

4 Quite a bit concerned

5 Greatly concerned

5. How often do you get your blood pressure checked?

1 Never or rarely

2 Once every 4 or 5 years

3 Once every 2 or 3 years

4 Once a year

5 More than once a year

6. About how many cigarettes do you smoke each day?

1 None

2 Less than 1/2 a pack

3 1/2 to 1 pack

4 1 to 2 packs

5 More than 2 packs

7. Having lung cancer is just a matter of fate so smoking doesn't make that big a difference. How much do you agree or disagree with this statement?

1 Strongly agree

2 Slightly agree

3 Do not agree or disagree

4 Slightly disagree

5 Strongly disagree

8. How concerned are you about being crippled by a heart attack?

1 Not concerned

2 Only a little concerned

3 Somewhat concerned

4 Quite a bit concerned

5 Greatly concerned

9. About how often do you exercise?

1 Not at all or hardly ever

2 Several times a year

3 One or two times a month

4 Two or three times a week

5 Daily or almost daily

10. How do you feel about the message you heard today? How much do you agree or disagree with this message?

1 Strongly agree

2 Slightly agree

3 Do not agree or disagree

4 Slightly disagree

5 Strongly disagree

11. Having a heart attack is just a matter of fate, so proper diet and exercise don't make that big a difference. How much do you agree or disagree with this statement?

1 Strongly agree

2 Slightly agree

3 Do not agree or disagree

4 Slightly disagree

5 Strongly disagree

12. Having a stroke is just a matter of fate so keeping my blood pressure down doesn't make that big a difference. How much do you agree or disagree with this statement?

1 Strongly agree

2 Slightly agree

3 Do not agree or disagree

4 Slightly disagree

5 Strongly disagree

13. How often do you go to have your eyes checked?

1 Less than once in three years

2 About once in three years

3 About once in two years

4 About once a year

5 I do not go unless my eyes give me trouble

14. The chances of having a stroke are so small that getting my blood pressure checked really isn't worth the trouble. How much do you agree or disagree with this statement?

1 Strongly agree

2 Slightly agree

3 Do not agree or disagree

4 Slightly disagree

5 Strongly disagree

15. How concerned are you about eating foods that will help keep you healthy?

1 Not at all concerned

2 Not very concerned

3 Not concerned one way or another

4 Somewhat concerned

5 Very concerned

16. How often do you go to a dentist to get your teeth checked?

1 Less than once in two years

2 About once in two years

3 About once a year

4 About once every six months

5 I do not go to the dentist except when my teeth bother me.

17. How concerned are you about being killed by a stroke?

1 Not at all concerned

2 Only a little concerned

3 Somewhat concerned

4 Quite a bit concerned

5 Greatly concerned

18. How concerned are you about getting lung cancer?

1 Not at all concerned

2 Only a little concerned

3 Somewhat concerned

4 Quite a bit concerned

5 Greatly concerned

19. Can you remember any other health or safety messages you have heard during the last week?

1 No

2 Yes Please describe the type of message heard: _____

20. In your opinion, what do you think are the three most important health problems in the U.S. today?

1. _____

2. _____

3. _____

What is your name? _____

What is your address? _____

What is your home telephone number? _____

How old are you? _____

What is your sex? Male _____ Female _____

What is your race? White _____ Black _____ Other _____

How far did you go in school?

_____ Less than high school
_____ High school
_____ More than high school

HEALTH STUDY QUESTIONNAIRE - 6

FOR RESEARCH ONLY - ALL YOUR ANSWERS WILL BE COMPLETELY CONFIDENTIAL

Please mark only one answer to each of the following questions:

1. How concerned are you about dying from a heart attack?

☐ Not concerned☐ Only a little concerned☐ Somewhat concerned☐ Quite a bit concerned☐ Greatly concerned

2. How often do you visit your doctor for a physical checkup?

☐ Less than once in three years☐ About once a year☐ About once in three years☐ More than once a year☐ About once in two years

3. How often do you use dental floss on your teeth?

☐ Never☐ A few times a month☐ Two or three times a week☐ Once a day☐ More than once a day

4. How concerned are you about being crippled by a stroke?

1 Not at all concerned

2 Only a little concerned

3 Somewhat concerned

4 Quite a bit concerned

5 Greatly concerned

5. How often do you get your blood pressure checked?

1 Never or rarely

2 Once every 4 or 5 years

3 Once every 2 or 3 years

4 Once a year

5 More than once a year

6. About how many cigarettes do you smoke each day?

1 None

2 Less than 1/2 a pack

3 1/2 to 1 pack

4 1 to 2 packs

5 More than 2 packs

7. Having lung cancer is just a matter of fate so smoking doesn't make that big a difference. How much do you agree or disagree with this statement?

1 Strongly agree

2 Slightly agree

3 Do not agree or disagree

4 Slightly disagree

5 Strongly disagree

8. How concerned are you about being crippled by a heart attack?

1 Not concerned

2 Only a little concerned

3 Somewhat concerned

4 Quite a bit concerned

5 Greatly concerned

9. About how often do you exercise?

1 Not at all or hardly ever

2 Several times a year

3 One or two times a month

4 Two or three times a week

5 Daily or almost daily

10. Having a heart attack is just a matter of fate, so proper diet and exercise don't make that big a difference. How much do you agree or disagree with this statement?

1 Strongly agree

2 Slightly agree

3 Do not agree or disagree

4 Slightly disagree

5 Strongly disagree

11. Having a stroke is just a matter of fate so keeping my blood pressure down doesn't make that big a difference. How much do you agree or disagree with this statement?

1 Strongly agree

2 Slightly agree

3 Do not agree or disagree

4 Slightly disagree

5 Strongly disagree

12. How often do you go to have your eyes checked?

1 Less than once in three years

2 About once in three years

3 About once in two years

4 About once a year

5 I do not go unless my eyes give me trouble

13. The chances of having a stroke are so small that getting my blood pressure checked really isn't worth the trouble. How much do you agree or disagree with this statement?

1 Strongly agree

2 Slightly agree

3 Do not agree or disagree

4 Slightly disagree

5 Strongly disagree

14. How concerned are you about eating foods that will help keep you healthy?

1 Not at all concerned

2 Not very concerned

3 Not concerned one way or another

4 Somewhat concerned

5 Very concerned

15. How often do you go to a dentist to get your teeth checked?

1 Less than once in two years

2 About once in two years

3 About once a year

4 About once every six months

5 I do not go to the dentist except when my teeth bother me.

16. How concerned are you about being killed by a stroke?

1 Not at all concerned

2 Only a little concerned

3 Somewhat concerned

4 Quite a bit concerned

5 Greatly concerned

17. How concerned are you about getting lung cancer?

1 Not at all concerned

2 Only a little concerned

3 Somewhat concerned

4 Quite a bit concerned

5 Greatly concerned

18. Can you remember any health or safety messages you have heard during the last week?

1 No

2 Yes Please describe the type of message heard: _____

19. In your opinion, what do you think are the three most important health problems in the U.S. today?

1. _____

2. _____

3.

What is your name? _____

What is your address? _____

What is your home telephone number? _____

How old are you? _____

What is your sex? Male _____ Female _____

What is your race? White _____ Black _____ Other _____

How far did you go in school?

_____ Less than high school
_____ High school
_____ More than high school

;

[illegible]

INSTRUCTIONS FOR MAKING TELEPHONE CALLS

Beginning the week of May 30 you are to call the telephone number shown below and listen to a 30-second recorded message. At the end of the message, leave your name and the date.

You must call on five (5) separate days each week for three weeks. If you can, please call on Monday, Tuesday, Wednesday, Thursday, and Friday, but if for some reason you miss a day, you may call in on Saturday or Sunday to make it up. You can call at any time of the day or night.

If you are out of town, you may call collect.

You must make all your telephone calls and then come in for your last meeting in order to complete the project.

Your special telephone number to call for your message is:

962-3463 (or 962-FINE)

Call five days between May 30 and June 5

Call five days between June 6 and June 12

Call five days between June 13 and June 19

If you need to make any changes in your last appointment, please call 962-6578 to work out the arrangements.

Thank you.

DO NOT LOSE THIS SHEET!

INSTRUCTIONS FOR MAKING TELEPHONE CALLS

Beginning the week of May 30 you are to call the telephone number shown below and listen to a 30-second recorded message. At the end of the message, leave your name and the date.

You must call on five (5) separate days each week for three weeks. If you can, please call on Monday, Tuesday, Wednesday, Thursday, and Friday, but if for some reason you miss a day, you may call in on Saturday or Sunday to make it up. You can call at any time of the day or night.

If you are out of town, you may call collect.

You must make all your telephone calls and then come in for your last meeting in order to complete the project.

Your special telephone number to call for your message is:

962-3438 (or 962-DIET)

Call five days between May 30 and June 5

Call five days between June 6 and June 12

Call five days between June 13 and June 19

If you need to make any changes in your last appointment, please call 962-6578 to work out the arrangements.

Thank you.

DO NOT LOSE THIS SHEET!

INSTRUCTIONS FOR MAKING TELEPHONE CALLS

Beginning the week of May 30 you are to call the telephone number shown below and listen to a 30-second recorded message. At the end of the message, leave your name and the date.

You must call on five (5) separate days each week for three weeks. If you can, please call on Monday, Tuesday, Wednesday, Thursday, and Friday, but if for some reason you miss a day, you may call in on Saturday or Sunday to make it up. You can call at any time of the day or night.

If you are out of town, you may call collect.

You must make all your telephone calls and then come in for your last meeting in order to complete the project.

Your special telephone number to call for your message is:

962-3463 (or 962-FINE)

Call five days between May 30 and June 5

Call five days between June 6 and June 12

Call five days between June 13 and June 19

If you need to make any changes in your last appointment, please call 962-6578 to work out the arrangements.

Thank you.

DO NOT LOSE THIS SHEET!

INSTRUCTIONS FOR MAKING TELEPHONE CALLS

Beginning the week of June 6 you are to call the telephone number shown below and listen to a 30-second recorded message. At the end of the message, leave your name and the date.

You must call on five (5) separate days each week for three weeks. If you can, please call on Monday, Tuesday, Wednesday, Thursday, and Friday, but if for some reason you miss a day, you may call in on Saturday or Sunday to make it up. You can call at any time of the day or night.

If you are out of town, you may call collect.

You must make all your telephone calls and then come in for your last meeting in order to complete the project.

Your special telephone number to call for your message is:

962-4278 (or 962-HART)

Call five days between June 6 and June 12

Call five days between June 13 and June 19

Call five days between June 20 and June 26

If you need to make any changes in your last appointment, please call 962-6578 to work out the arrangements.

Thank you.

DO NOT LOSE THIS SHEET!

INSTRUCTIONS FOR MAKING TELEPHONE CALLS

Beginning the week of June 6 you are to call the telephone number shown below and listen to a 30-second recorded message. At the end of the message, leave your name and the date.

You must call on five (5) separate days each week for three weeks. If you can, please call on Monday, Tuesday, Wednesday, Thursday, and Friday, but if for some reason you miss a day, you may call in on Saturday or Sunday to make it up. You can call at any time of the day or night.

If you are out of town, you may call collect.

You must make all your telephone calls and then come in for your last meeting in order to complete the project.

Your special telephone number to call for your message is:

962-4278 (or 962-HART)

Call five days between June 6 and June 12

Call five days between June 13 and June 19

Call five days between June 20 and June 26

If you need to make any changes in your last appointment, please call 962-6578 to work out the arrangements.

Thank you.

DO NOT LOSE THIS SHEET!

Appendix 10. Information Form Explaining Incentive Raffle

Health Research Study

The drawing for the Microwave Oven and additional prizes will be held on July 15.

The winners will be contacted by telephone, and informed when and where to pick up their prizes.

Anyone interested in knowing who won the lottery prizes may call 966-1044 from July 20 - July 31. A list of winners will be given by a recorded message.

The prizes are: 1 Grand Prize: A Microwave Oven

3 Second Prizes: \$30.00 Gift Certificate
from Belk

8 Third Prizes: \$20.00 Gift Certificate
from Belk

12 Fourth Prizes: \$10.00 Gift Certificate
from Belk

Appendix 11. Tables of Preliminary Cross-Tabs of Questionnaire Responses, Subject Demographics, Changes in Belt Usage, and Message Groups

I. Crosstabs of belt wearing (INBELT1) with questionnaire responses (from first meeting) concerning other health maintenance behaviors for all groups .

1. INBELT1 By Quest2 (Checkup Frequency)	(N.S.)
2. INBELT1 By Quest3 (Flossing Frequency)	(p <.03)
3. INBELT1 By Quest6 (Smoking)	(N.S.)
4. INBELT1 By Quest9 (Exercise Frequency)	(N.S.)
5. INBELT1 By Quest12 (Eye Checkup Frequency)	(p <.10)
6. INBELT1 By Quest15 (Dental Checkup Frequency)	(N.S.)
7. INBELT1 By Race	(N.S.)
8. INBELT1 By Sex	(N.S.)
9. INBELT1 By Age	(N.S.)

II. Crosstabs of seat belt question (Quest5) with responses to other health maintenance activities on the questionnaire for Groups 1 to 5 .

1. Quest5 By Quest2 (Checkup Frequency)	(N.S.)
2. Quest5 By Quest3 (Flossing Frequency)	(N.S.)
3. Quest5 By Quest6 (Smoking)	(N.S.)
4. Quest5 By Quest9 (Exercise)	(N.S.)
5. Quest5 By Quest12 (Eye Checkup Frequency)	(p <.005)
6. Quest5 By Quest15 (Dental Checkup Frequency)	(N.S.)
7. INBELT1 By Quest5	(p <.0001)

III. Crosstabs of change in belt wearing and in questionnaire responses by Group and demographic variables for all Groups .

1. Group By N1 (Question on Heart Attack)	(N.S.)
2. Group By N2 (Question on Checkup Frequency)	(N.S.)
3. Group By N3 (Question on Flossing Frequency)	(N.S.)
4. Group By N4 (Question on Being Crippled by Accident/Stroke)	(p=.0006)
5. Group By N5 (Question on Wearing Seat Belts/Checking Blood Pressure)	(p=.0015)
6. Group By N6 (Question on Smoking)	(N.S.)
7. Group By N7 (Question on Smoking and Lung Cancer)	(N.S.)
8. Group By N8 (Question on Being Crippled by Heart Attack)	(N.S.)
9. Group By N9 (Question on Exercise Frequency)	(N.S.)
10. Group By N10 (Question on Diet and Exercise on Heart)	(N.S.)
11. Group By N11 (Question on Belt/Blood Pressure)	(N.S.)
12. Group By N12 (Question on Eye Checkup Frequency)	(N.S.)
13. Group By N13 (Question on Belt/Blood Pressure)	(N.S.)
14. Group By N14 (Question on Eating Habits)	(N.S.)
15. Group By N15 (Question on Dental Checkup Frequency)	(N.S.)
16. Group By N16 (Question on Being Killed By Accident/Stroke)	(N.S.)
17. Group By N17 (Question on Concern for Lung Cancer)	(N.S.)
18. Group By R (Change in Seat Belt Wearing)	(N.S.)
19. R (Change in Seat Belt Wearing) By Race	(N.S.)
20. R (Change in Seat Belt Wearing) By Sex	(N.S.)
21. R (Change in Seat Belt Wearing) By Age	(N.S.)

IV. Crosstabs of Change in Seat Belt Wearing and in Response to Seat Belt Questions by Demographic Variables and by Changes in response to selected Questionnaire Items for Groups 1-5 .

1.	N5 (Wearing Seat Belt/Checking Blood Pressure) By Race	(p=.0002)
2.	N5 (Wearing Seat Belt/Checking Blood Pressure) By Sex	(N.S.)
3.	N5 (Wearing Seat Belt/Checking Blood Pressure) By Age	(N.S.)
4.	N5 (Wearing Seat Belt/Checking Blood Pressure) By R (Change in Seat Belt Wearing)	(N.S.)
5.	N5 (Wearing Seat Belt/Checking Blood Pressure) By N2 (Checkup Frequency)	(p=.015)
6.	N5 (Wearing Seat Belt/Checking Blood Pressure) By N6 (Smoking)	(N.S.)
7.	N11 (Belt/Blood Pressure) By Race	(N.S.)
8.	N11 (Belt/Blood Pressure) By Sex	(N.S.)
9.	N11 (Belt/Blood Pressure) By Age	(N.S.)
10.	N11 (Belt/Blood Pressure) By R (Change in Seat Belt Wearing)	(N.S.)
11.	N11 (Belt/Blood Pressure) By N2 (Checkup Frequency)	(N.S.)
12.	N11 (Belt/Blood Pressure) By N6 (Smoking)	(N.S.)
13.	N16 (Being Killed by Accident/Stroke) By Race	(p=.029)
14.	N16 (Being Killed by Accident/Stroke) By Sex	(p=.0089)
15.	N16 (Being killed by Accident/Stroke) By Age	(N.S.)
16.	N16 (Being Killed by Accident/Stroke) By R (Change in Seat Belt Wearing)	(N.S.)
17.	N16 (Being Killed by Accident/Stroke) By N2 (Checkup Frequency)	(p=.0002)
18.	N16 (Being Killed by Accident/Stroke) By N6 (Smoking)	(N.S.)

ALL GROUPS

TABLE OF INBELT1 BY QUEST2

INBELT1	QUEST2					CHECK UP
FREQUENCY						
PERCENT						
ROW PCT						
COL PCT	1	2	3	4	5	TOTAL
B	16	16	17	37	6	92
	7.11	7.11	7.56	16.44	2.67	40.89
	17.39	17.39	18.48	40.22	6.52	
	57.14	50.00	38.64	37.37	27.27	
N	12	16	27	62	16	133
	5.33	7.11	12.00	27.56	7.11	59.11
	9.02	12.03	20.30	46.62	12.03	
	42.86	50.00	61.36	62.63	72.73	
TOTAL	28	32	44	99	22	225
	12.44	14.22	19.56	44.00	9.78	100.00

CHI-SQUARE 6.446 DF= 4 PROB=0.1682
 PHI 0.169
 CONTINGENCY COEFFICIENT 0.167
 CRAMER'S V 0.169
 LIKELIHOOD RATIO CHISQUARE 6.452 DF= 4 PROB=0.1679

ALL GROUPS

TABLE OF INBELT1 BY QUEST3

INBELT1	QUEST3					FLOSSING
FREQUENCY						
PERCENT						
ROW PCT						
COL PCT	1	2	3	4	5	TOTAL
B	5	33	31	23	0	92
	2.22	14.67	13.78	10.22	0.00	40.89
	5.43	35.87	33.70	25.00	0.00	
	19.23	40.74	43.66	53.49	0.00	
N	21	48	40	20	4	133
	9.33	21.33	17.78	8.89	1.78	59.11
	15.79	36.09	30.08	15.04	3.01	
	80.77	59.26	56.34	46.51	100.00	
TOTAL	26	81	71	43	4	225
	11.56	36.00	31.56	19.11	1.78	100.00

CHI-SQUARE 10.864 DF= 4 PROB=0.0261
 PHI 0.220
 CONTINGENCY COEFFICIENT 0.215
 CRAMER'S V 0.220
 LIKELIHOOD RATIO CHISQUARE 12.766 DF= 4 PROB=0.0125

ALL GROUPS

TABLE OF INBELT1 BY QUEST6

INBELT1	QUEST6		SMOKING				
FREQUENCY	PERCENT						
ROW PCT	COL PCT						
	0	1	2	3	4	5	TOTAL
F	0	1	5	2	4	80	92
	0.00	0.44	2.22	0.89	1.78	35.56	40.89
	0.00	1.09	5.43	2.17	4.35	86.96	
	0.00	100.00	41.67	15.38	23.53	44.20	
N	1	0	7	11	13	101	133
	0.44	0.00	3.11	4.89	5.78	44.89	59.11
	0.75	0.00	5.26	8.27	9.77	75.94	
	100.00	0.00	58.33	84.62	76.47	55.80	
TOTAL	1	1	12	13	17	181	225
	0.44	0.44	5.33	5.78	7.56	80.44	100.00

CHI-SQUARE	8.579	DF=	5	PROB=0.1271
PHI	0.195			
CONTINGENCY COEFFICIENT	0.192			
CRAMER'S V	0.195			
LIKELIHOOD RATIO CHISQUARE	9.913	DF=	5	PROB=0.0777

ALL GROUPS

TABLE OF INBELT1 BY QUEST9

INBELT1	QUEST9		EXERCISE							
FREQUENCY	PERCENT	ROW PCT	COL PCT	0	1	2	3	4	5	TOTAL
-----+-----+-----+-----+-----+-----+-----+-----+-----+-----+-----										
E	0	5	3	15	34	35				92
	0.00	2.22	1.33	6.67	15.11	15.56				40.89
	0.00	5.43	3.26	16.30	36.96	38.04				
	0.00	27.78	27.27	55.56	38.64	43.75				
-----+-----+-----+-----+-----+-----+-----+-----+-----+-----+-----										
N	1	13	8	12	54	45				133
	0.44	5.78	3.56	5.33	24.00	20.00				59.11
	0.75	9.77	6.02	9.02	40.60	33.83				
	100.00	72.22	72.73	44.44	61.36	56.25				
-----+-----+-----+-----+-----+-----+-----+-----+-----+-----+-----										
TOTAL	1	18	11	27	88	80				225
	0.44	8.00	4.89	12.00	39.11	35.56				100.00

CHI-SQUARE	5.674	DF=	5	PROB=0.3392
PHI	0.159			
CONTINGENCY COEFFICIENT	0.157			
CRAMER'S V	0.159			
LIKELIHOOD RATIO CHISQUARE	6.087	DF=	5	PROB=0.2978

ALL GROUPS

TABLE OF INBELT1 BY QUEST12

INBELT1 QUEST12 EYES CHECKED

FREQUENCY						
PERCENT						
ROW PCT						
COL PCT	1	2	3	4	5	TOTAL
R	25	6	22	25	14	92
	11.16	2.68	9.82	11.16	6.25	41.07
	27.17	6.52	23.91	27.17	15.22	
	43.86	30.00	53.66	45.45	27.45	
N	32	14	19	30	37	132
	14.29	6.25	8.48	13.39	16.52	58.93
	24.24	10.61	14.39	22.73	28.03	
	56.14	70.00	46.34	54.55	72.55	
TOTAL	57	20	41	55	51	224
	25.45	8.93	18.30	24.55	22.77	100.00

CHI-SQUARE	8.226	DF=	4	PROB=0.0837
PHI	0.192			
CONTINGENCY COEFFICIENT	0.188			
CRAMER'S V	0.192			
LIKELIHOOD RATIO CHISQUARE	8.403	DF=	4	PROB=0.0779

ALL GROUPS

TABLE OF INBELT1 BY QUEST15

INBELT1 QUEST15 TEETH CHECKED

FREQUENCY						
PERCENT						
ROW PCT						
COL PCT	1	2	3	4	5	TOTAL
B	2	18	30	35	7	92
	0.89	8.00	13.33	15.56	3.11	40.89
	2.17	19.57	32.61	38.04	7.61	
	15.38	45.00	39.47	43.75	43.75	
N	11	22	46	45	9	133
	4.89	9.78	20.44	20.00	4.00	59.11
	8.27	16.54	34.59	33.83	6.77	
	84.62	55.00	60.53	56.25	56.25	
TOTAL	13	40	76	80	16	225
	5.78	17.78	33.78	35.56	7.11	100.00

CHI-SQUARE	4.166	DF=	4	PROB=0.3840
PHI	0.136			
CONTINGENCY COEFFICIENT	0.135			
CRAMER'S V	0.136			
LIKELIHOOD RATIO CHISQUARE	4.645	DF=	4	PROB=0.3257

ALL GROUPS

TABLE OF INBELT1 BY RACE

INBELT1	RACE		
FREQUENCY			
PERCENT			
ROW PCT			
COL PCT	A	W	TOTAL
B	7	85	92
	3.11	37.78	40.89
	7.61	92.39	
	28.00	42.50	
N	18	115	133
	8.00	51.11	59.11
	13.53	86.47	
	72.00	57.50	
TOTAL	25	200	225
	11.11	88.89	100.00

CHI-SQUARE	1.933	DF=	1	PROB=0.1644
PHI	-0.093			
CONTINGENCY COEFFICIENT	0.092			
CRAMER'S V	0.093			
LIKELIHOOD RATIO CHISQUARE	2.014	DF=	1	PROB=0.1559
CONTINUITY ADJ. CHI-SQUARE	1.380	DF=	1	PROB=0.2402
FISHER'S EXACT TEST (1-TAIL)				PROB=0.1190
(2-TAIL)				PROB=0.1986

ALL GROUPS

TABLE OF INBELT4 BY SEX

INBELT1	SEX		
FREQUENCY			
PERCENT			
ROW PCT			
COL PCT	F	M	TOTAL
-----+-----+-----+-----			
E	73	19	92
	32.44	8.44	40.89
	79.35	20.65	
	42.69	35.19	
-----+-----+-----+-----			
N	98	35	133
	43.56	15.56	59.11
	73.68	26.32	
	57.31	64.81	
-----+-----+-----+-----			
TOTAL	171	54	225
	76.00	24.00	100.00

CHI-SQUARE	0.956	DF=	1	PROB=0.3281
PHI	0.065			
CONTINGENCY COEFFICIENT	0.065			
CRAMER'S V	0.065			
LIKELIHOOD RATIO CHISQUARE	0.968	DF=	1	PROB=0.3252
CONTINUITY ADJ. CHI-SQUARE	0.671	DF=	1	PROB=0.4127
FISHER'S EXACT TEST (1-TAIL)				PROB=0.2069
(2-TAIL)				PROB=0.3458

ALL GROUPS

TABLE OF INBELT1 BY AGE

INBELT1	AGE			
FREQUENCY				
PERCENT				
ROW PCT				
COL PCT	18-25	26-35	36+	TOTAL
F	23	42	27	92
	10.22	18.67	12.00	40.89
	25.00	45.65	29.35	
	34.33	49.41	36.99	
N	44	43	46	133
	19.56	19.11	20.44	59.11
	33.08	32.33	34.59	
	65.67	50.59	63.01	
TOTAL	67	85	73	225
	29.78	37.78	32.44	100.00

CHI-SQUARE 4.208 DF= 2 PROB=0.1220
 PHI 0.137
 CONTINGENCY COEFFICIENT 0.135
 CRAMER'S V 0.137
 LIKELIHOOD RATIO CHISQUARE 4.196 DF= 2 PROB=0.1227

GROUPS 1-5

TABLE OF QUEST5 BY QUEST2

QUEST5	QUEST2					CHECK UP
FREQUENCY PERCENT ROW PCT COL PCT	1	2	3	4	5	TOTAL
1	3 1.64 13.64 13.04	2 1.09 9.09 8.33	7 3.63 31.82 17.95	8 4.37 36.36 10.26	2 1.09 9.09 10.53	22 12.02
2	3 1.64 10.71 13.04	4 2.19 14.29 16.67	5 2.73 17.86 12.82	13 7.10 46.43 16.67	3 1.64 10.71 15.79	28 15.30
3	3 1.64 7.69 13.04	2 1.09 5.13 8.33	11 6.01 28.21 28.21	17 9.29 43.59 21.79	6 3.26 15.38 31.58	39 21.31
4	5 2.73 12.50 21.74	6 3.28 15.00 25.00	9 4.92 22.50 23.08	16 8.74 40.00 20.51	4 2.19 10.00 21.05	40 21.66
5	9 4.92 16.67 39.13	10 5.46 18.52 41.67	7 3.83 12.96 17.95	24 13.11 44.44 30.77	4 2.19 7.41 21.05	54 29.51
TOTAL	23 12.57	24 13.11	39 21.31	78 42.62	19 10.38	183 100.00

CHI-SQUARE 10.680 DF= 16 PROB=0.6168
 PHI 0.244
 CONTINGENCY COEFFICIENT 0.237
 CRAMER'S V 0.122
 LIKELIHOOD RATIO CHISQUARE 11.408 DF= 16 PROB=0.7636

GROUPS 1-5

TABLE OF QUEST5 BY QUEST3

QUEST5	QUEST3					FLOSSING
FREQUENCY						
PERCENT						
ROW PCT						
COL PCT	1	2	3	4	5	TOTAL
1	5	7	6	2	2	22
	2.73	3.83	3.28	1.09	1.09	12.02
	22.73	31.82	27.27	9.09	9.09	
	21.74	10.00	12.00	5.56	50.00	
2	7	8	6	7	0	28
	3.83	4.37	3.28	3.83	0.00	15.30
	25.00	28.57	21.43	25.00	0.00	
	30.43	11.43	12.00	19.44	0.00	
3	5	15	12	6	1	39
	2.73	8.20	6.56	3.28	0.55	21.31
	12.82	38.46	30.77	15.38	2.56	
	21.74	21.43	24.00	16.67	25.00	
4	3	20	10	7	0	40
	1.64	10.93	5.46	3.83	0.00	21.86
	7.50	50.00	25.00	17.50	0.00	
	13.04	28.57	20.00	19.44	0.00	
5	3	20	16	14	1	54
	1.64	10.93	8.74	7.65	0.55	29.51
	5.56	37.04	29.63	25.93	1.85	
	13.04	28.57	32.00	38.89	25.00	
TOTAL	23	70	50	36	4	183
	12.57	38.25	27.32	19.67	2.19	100.00

CHI-SQUARE 20.609 DF= 16 PROB=0.1860
 PHI 0.337
 CONTINGENCY COEFFICIENT 0.320
 CRAMER'S V 0.169
 LIKELIHOOD RATIO CHISQUARE 19.925 DF= 16 PROB=0.2236

GROUPS 1-5

TABLE OF QUEST5 BY QUEST6

QUEST5	QUEST6	SMOKING					
FREQUENCY PERCENT ROW PCT COL PCT	0	1	2	3	4	5	TOTAL
1	1 0.55 4.55 100.00	0 0.00 0.00 0.00	1 0.55 4.55 9.09	4 2.19 16.18 33.33	0 0.00 0.00 0.00	16 8.74 72.73 11.27	22 12.02
2	0 0.00 0.00 0.00	0 0.00 0.00 0.00	1 0.55 3.57 9.09	1 0.55 3.57 8.33	3 1.64 10.71 18.75	23 12.57 82.14 16.20	28 15.30
3	0 0.00 0.00 0.00	0 0.00 0.00 0.00	2 1.09 5.13 18.18	4 2.19 16.26 33.33	7 3.83 17.95 43.75	26 14.21 66.67 18.31	39 21.31
4	0 0.00 0.00 0.00	0 0.00 0.00 0.00	2 1.09 5.00 18.18	2 1.09 5.00 16.67	4 2.19 10.00 25.00	32 17.49 80.00 22.54	40 21.86
5	0 0.00 0.00 0.00	1 0.55 1.85 100.00	5 2.73 9.26 45.45	1 0.55 1.85 8.33	2 1.09 3.70 12.50	45 24.59 83.33 31.69	54 29.51
TOTAL	1 0.55	1 0.55	11 6.01	12 6.56	16 8.74	142 77.60	183 100.00

CHI-SQUARE 27.299 DF= 20 PROB=0.1271
 PHI 0.386
 CONTINGENCY COEFFICIENT 0.360
 CRAMER'S V 0.193
 LIKELIHOOD RATIO CHISQUARE 24.889 DF= 20 PROB=0.2057

GROUPS 1-5

TABLE OF QUEST5 BY QUEST9

QUEST5	QUEST9	EXERCISE					
FREQUENCY							
PERCENT							
ROW PCT							
COL PCT	0	1	2	3	4	5	TOTAL
1	0	4	2	2	8	6	22
	0.00	2.19	1.09	1.09	4.37	3.28	12.02
	0.00	18.18	9.09	9.09	36.36	27.27	
	0.00	26.67	22.22	6.00	11.94	9.09	
2	1	2	2	3	6	14	28
	0.55	1.09	1.09	1.64	3.28	7.65	15.30
	3.57	7.14	7.14	10.71	21.43	50.00	
	100.00	13.33	22.22	12.00	8.96	21.21	
3	0	4	3	5	14	13	39
	0.00	2.19	1.64	2.73	7.65	7.10	21.31
	0.00	10.26	7.69	12.82	35.90	33.33	
	0.00	26.67	33.33	20.00	20.90	19.70	
4	0	2	0	8	16	14	40
	0.00	1.09	0.00	4.37	8.74	7.65	21.86
	0.00	5.00	0.00	20.00	40.00	35.00	
	0.00	13.33	0.00	32.00	23.88	21.21	
5	0	3	2	7	23	19	54
	0.00	1.64	1.09	3.83	12.57	10.38	29.51
	0.00	5.56	3.70	12.96	42.59	35.19	
	0.00	20.00	22.22	26.00	34.33	28.79	
TOTAL	1	15	9	25	67	66	183
	0.55	8.20	4.92	13.66	36.61	36.07	100.00

CHI-SQUARE 19.446 DF= 20 PROB=0.4930
 PHI 0.326
 CONTINGENCY COEFFICIENT 0.310
 CRAMER'S V 0.163
 LIKELIHOOD RATIO CHISQUARE 18.672 DF= 20 PROB=0.5302

GROUPS 1-5

TABLE OF QUEST5 BY QUEST12

QUEST5	QUEST12					EYES CHECKED
FREQUENCY						
PERCENT						
ROW PCT						
COL PCT	1	2	3	4	5	TOTAL
1	6	7	1	2	6	22
	3.28	3.83	0.55	1.09	3.28	12.02
	27.27	31.82	4.55	9.09	27.27	
	12.50	41.18	3.33	4.76	13.04	
2	8	0	3	7	10	28
	4.37	0.00	1.64	3.83	5.46	15.30
	28.57	0.00	10.71	25.00	35.71	
	16.67	0.00	10.00	16.67	21.74	
3	10	3	5	6	15	39
	5.46	1.64	2.73	3.28	8.20	21.31
	25.64	7.69	12.82	15.38	38.46	
	20.83	17.65	16.67	14.29	32.61	
4	7	4	8	14	7	40
	3.83	2.19	4.37	7.65	3.83	21.86
	17.50	10.00	20.00	35.00	17.50	
	14.58	23.53	26.67	33.33	15.22	
5	17	3	13	13	9	54
	9.29	1.64	7.10	7.10	4.37	29.51
	31.48	5.56	24.07	24.07	14.81	
	35.42	17.65	43.33	30.95	17.39	
TOTAL	48	17	30	42	46	183
	26.23	9.29	16.39	22.95	25.14	100.00

CHI-SQUARE 35.045 DF= 16 PROB=0.0039
 PHI 0.438
 CONTINGENCY COEFFICIENT 0.401
 CRAMER'S V 0.219
 LIKELIHOOD RATIO CHISQUARE 34.114 DF= 16 PROB=0.0052

GROUPS 1-5

TABLE OF QUEST5 BY QUEST15

QUEST5	QUEST15					TEETH CHECKED
FREQUENCY						
PERCENT						
ROW PCT						
COL PCT	1	2	3	4	5	TOTAL
1	3	4	4	9	2	22
	1.64	2.19	2.19	4.92	1.09	12.02
	13.64	18.18	18.18	40.91	9.09	
	30.00	12.90	6.35	13.43	16.67	
2	1	4	7	14	2	28
	0.55	2.19	3.83	7.65	1.09	15.30
	3.57	14.29	25.00	50.00	7.14	
	10.00	12.90	11.11	20.90	16.67	
3	4	5	16	12	2	39
	2.19	2.73	8.74	6.56	1.09	21.31
	10.26	12.82	41.03	30.77	5.13	
	40.00	16.13	25.40	17.91	16.67	
4	2	8	17	12	1	40
	1.09	4.37	9.29	6.56	0.55	21.86
	5.00	20.00	42.50	30.00	2.50	
	20.00	25.81	26.98	17.91	8.33	
5	0	10	19	20	5	54
	0.00	5.46	10.38	10.93	2.73	29.51
	0.00	18.52	35.19	37.04	9.26	
	0.00	32.26	30.16	29.85	41.67	
TOTAL	10	31	63	67	12	183
	5.46	16.94	34.43	36.61	6.56	100.00

CHI-SQUARE 16.257 DF= 16 PROB=0.4352
 PHI 0.298
 CONTINGENCY COEFFICIENT 0.286
 CRAMER'S V 0.149
 LIKELIHOOD RATIO CHISQUARE 18.611 DF= 16 PROB=0.2894

GROUPS 1-5

TABLE OF INBELT1 BY QUEST5

INBELT1	QUEST5					
FREQUENCY						
PERCENT						
ROW PCT						
COL PCT	1	2	3	4	5	TOTAL
E	0	2	4	23	49	78
	0.00	1.10	2.20	12.64	26.92	42.86
	0.00	2.56	5.13	29.49	62.82	
	0.00	7.14	10.53	57.50	90.74	
N	22	26	34	17	5	104
	12.09	14.29	18.68	9.34	2.75	57.14
	21.15	25.00	32.69	16.35	4.81	
	100.00	92.86	89.47	42.50	9.26	
TOTAL	22	28	38	40	54	182
	12.09	15.38	20.86	21.98	29.67	100.00

CHI-SQUAPE	101.562	DF=	4	PROB=0.0001
PHI	0.746			
CONTINGENCY COEFFICIENT	0.598			
CRAMER'S V	0.746			
LIKELIHOOD RATIO CHISQUARE	120.729	DF=	4	PROB=0.0001

ALL GROUPS

TABLE OF GROUP BY N1

GROUP	FINAL GROUP	N1	HEART ATTACK	
FREQUENCY	PERCENT	ROW PCT	COL PCT	TOTAL
	HIGH	MEDIUM	LOW	
1	2	7	30	39
	0.86	3.10	13.27	17.26
	5.13	17.95	76.92	
	9.52	28.00	16.67	
2	3	8	27	38
	1.33	3.54	11.95	16.81
	7.89	21.05	71.05	
	14.29	32.00	15.00	
3	3	5	32	40
	1.33	2.21	14.16	17.70
	7.50	12.50	80.00	
	14.29	20.00	17.76	
4	6	2	25	33
	2.65	0.88	11.06	14.60
	18.18	6.06	75.76	
	28.57	8.00	13.89	
5	4	2	27	33
	1.77	0.88	11.95	14.60
	12.12	6.06	81.82	
	19.05	8.00	15.00	
6	3	1	39	43
	1.33	0.44	17.26	19.03
	6.98	2.33	90.70	
	14.29	4.00	21.67	
TOTAL	21	25	180	226
	9.29	11.06	79.65	100.00

CHI-SQUARE 15.051 DF= 10 PROB=0.1302
 PHI 0.258
 CONTINGENCY COEFFICIENT 0.250
 CRAMER'S V 0.182
 LIKELIHOOD RATIO CHISQUARE 15.300 DF= 10 PROB=0.1215

ALL GROUPS

TABLE OF GROUP PY N2

GROUP	FINAL GROUP	N2	CHECKUP	
FREQUENCY				
PERCENT				
ROW PCT				
COL PCT	HIGH	MEDIUM	LOW	TOTAL
1	6	1	32	39
	2.65	0.44	14.16	17.26
	15.38	2.56	82.05	
	13.64	5.88	19.39	
2	9	2	27	38
	3.98	0.88	11.95	16.81
	23.68	5.26	71.05	
	20.45	11.76	16.36	
3	9	5	26	40
	3.98	2.21	11.50	17.70
	22.50	12.50	65.00	
	20.45	29.41	15.76	
4	6	1	26	33
	2.65	0.44	11.50	14.60
	18.18	3.03	78.79	
	13.64	5.88	15.76	
5	7	3	23	33
	3.10	1.33	10.18	14.60
	21.21	9.09	69.70	
	15.91	17.65	13.94	
6	7	5	31	43
	3.10	2.21	13.72	19.03
	16.28	11.63	72.09	
	15.91	29.41	18.79	
TOTAL	44	17	165	226
	19.47	7.52	73.01	100.00

CHI-SQUARE	6.994	DF=	10	PROB=0.7260
PHI	0.176			
CONTINGENCY COEFFICIENT	0.173			
CRAMER'S V	0.124			
LIKELIHOOD RATIO CHISQUARE	7.360	DF=	10	PROB=0.6911

ALL GROUPS

TABLE OF GROUP BY N3

GROUP	FINAL GROUP	N3	FLOSS	
FREQUENCY				
PERCENT				
ROW PCT				
COL PCT	HIGH	MEDIUM	LOW	TOTAL
1	24	2	13	39
	10.62	0.88	5.75	17.26
	61.54	5.13	33.33	
	20.34	11.11	14.44	
2	18	3	17	38
	7.96	1.33	7.52	16.81
	47.37	7.89	44.74	
	15.25	16.67	18.89	
3	24	1	15	40
	10.62	0.44	6.64	17.70
	60.00	2.50	37.50	
	20.34	5.56	16.67	
4	15	4	14	33
	6.64	1.77	6.19	14.60
	45.45	12.12	42.42	
	12.71	22.22	15.56	
5	16	3	14	33
	7.08	1.33	6.19	14.60
	48.48	9.09	42.42	
	13.56	16.67	15.56	
6	21	5	17	43
	9.29	2.21	7.52	19.03
	48.84	11.63	39.53	
	17.80	27.78	18.89	
TOTAL	118	18	90	226
	52.21	7.96	39.82	100.00

CHI-SQUARE	5.952	DF=	10	PROB=0.8193
PHI	0.162			
CONTINGENCY COEFFICIENT	0.160			
CRAMER'S V	0.115			
LIKELIHOOD RATIO CHISQUARE	6.352	DF=	10	PROB=0.7849

ALL GROUPS

TABLE OF GROUP BY N4

GROUP	FINAL GROUP	N4	ACCIDENT/STR	
FREQUENCY				
PERCENT				
ROW PCT				
COL PCT	HIGH	MEDIUM	LOW	TOTAL
1	7	3	29	39
	3.10	1.33	12.83	17.26
	17.95	7.69	74.36	
	9.33	42.86	20.14	
2	17	2	19	38
	7.52	0.88	8.41	16.81
	44.74	5.26	50.00	
	22.67	28.57	13.19	
3	12	0	28	40
	5.31	0.00	12.39	17.70
	30.00	0.00	70.00	
	16.00	0.00	19.44	
4	10	0	23	33
	4.42	0.00	10.18	14.60
	30.30	0.00	69.70	
	13.33	0.00	15.97	
5	4	1	28	33
	1.77	0.44	12.39	14.60
	12.12	3.03	84.85	
	5.33	14.29	19.44	
6	25	1	17	43
	11.06	0.44	7.52	19.03
	58.14	2.33	39.53	
	33.33	14.29	11.81	
TOTAL	75	7	144	226
	33.19	3.10	63.72	100.00

CHI-SQUARE 31.628 DF= 10 PROB=0.0006
 PHI 0.371
 CONTINGENCY COEFFICIENT 0.347
 CRAMER'S V 0.262
 LIKELIHOOD RATIO CHISQUARE 33.203 DF= 10 PROB=0.0003

ALL GROUPS

TABLE OF GROUP BY N5

GROUP	FINAL GROUP	N5	BELT/BP	
FREQUENCY				
PERCENT				
ROW PCT				
COL PCT	HIGH	MEDIUM	LOW	TOTAL
1	14	4	21	39
	6.19	1.77	9.29	17.26
	35.90	10.26	53.85	
	28.57	33.33	12.73	
2	4	1	33	38
	1.77	0.44	14.60	16.81
	10.53	2.63	86.84	
	8.16	8.33	20.00	
3	6	4	30	40
	2.65	1.77	13.27	17.70
	15.00	10.00	75.00	
	12.24	33.33	18.18	
4	12	2	19	33
	5.31	0.88	8.41	14.60
	36.36	6.06	57.58	
	24.49	16.67	11.52	
5	10	1	22	33
	4.42	0.44	9.73	14.60
	30.30	3.03	66.67	
	20.41	8.33	13.33	
6	3	0	40	43
	1.33	0.00	17.70	19.03
	6.98	0.00	93.02	
	6.12	0.00	24.24	
TOTAL	49	12	165	226
	21.68	5.31	73.01	100.00

CHI-SQUARE	28.546	DF=	10	PROB=0.0015
PHI	0.355			
CONTINGENCY COEFFICIENT	0.335			
CRAMER'S V	0.251			
LIKELIHOOD RATIO CHISQUARE	31.192	DF=	10	PROB=0.0005

ALL GROUPS

TABLE OF GROUP BY N6

GROUP	FINAL GROUP	N6	SMOKING	
FREQUENCY				
PERCENT				
ROW PCT				
COL PCT	HIGH	MEDIUM	LOW	TOTAL
1	1	0	38	39
	0.44	0.00	16.81	17.26
	2.56	0.00	97.44	
	5.88	0.00	18.81	
2	3	1	34	38
	1.33	0.44	15.04	16.81
	7.89	2.63	89.47	
	17.65	14.29	16.83	
3	3	3	34	40
	1.33	1.33	15.04	17.70
	7.50	7.50	85.00	
	17.65	42.86	16.83	
4	5	1	27	33
	2.21	0.44	11.95	14.60
	15.15	3.03	81.82	
	29.41	14.29	13.37	
5	2	1	30	33
	0.88	0.44	13.27	14.60
	6.06	3.03	90.91	
	11.76	14.29	14.85	
6	3	1	39	43
	1.33	0.44	17.26	19.03
	6.98	2.33	90.70	
	17.65	14.29	19.31	
TOTAL	17	7	202	226
	7.52	3.10	89.38	100.00

CHI-SQUARE 8.364 DF= 10 PROB=0.5933
 PHI 0.192
 CONTINGENCY COEFFICIENT 0.189
 CRAMER'S V 0.136
 LIKELIHOOD RATIO CHISQUARE 8.723 DF= 10 PROB=0.5586

ALL GROUPS

TABLE OF GROUP BY N7

GROUP	FINAL GROUP	N7	LUNG CANCER	
FREQUENCY				
PERCENT				
ROW PCT				
COL PCT	HIGH	MEDIUM	LOW	TOTAL
1	6	0	33	39
	2.65	0.00	14.60	17.26
	15.38	0.00	84.62	
	21.43	0.00	17.01	
2	5	1	32	38
	2.21	0.44	14.16	16.81
	13.16	2.63	84.21	
	17.86	25.00	16.49	
3	5	1	34	40
	2.21	0.44	15.04	17.70
	12.50	2.50	85.00	
	17.86	25.00	17.53	
4	4	1	28	33
	1.77	0.44	12.39	14.60
	12.12	3.03	84.85	
	14.29	25.00	14.43	
5	5	0	28	33
	2.21	0.00	12.39	14.60
	15.15	0.00	84.85	
	17.86	0.00	14.43	
6	3	1	39	43
	1.33	0.44	17.26	19.03
	6.98	2.33	90.70	
	10.71	25.00	20.10	
TOTAL	28	4	194	226
	12.39	1.77	85.84	100.00

CHI-SQUARE 3.596 DF= 10 PROB=0.9637
 PHI 0.126
 CONTINGENCY COEFFICIENT 0.125
 CRAMER'S V 0.089
 LIKELIHOOD RATIO CHISQUARE 4.932 DF= 10 PROB=0.8957

ALL GROUPS

TABLE OF GROUP BY N8

GROUP	FINAL GROUP	N8	HEART ATTACK		
FREQUENCY					
PERCENT					
ROW PCT					
COL PCT	HIGH	MEDIUM	LOW		TOTAL
1	30	3	6		39
	13.27	1.33	2.65		17.26
	76.92	7.69	15.38		
	16.57	15.00	24.00		
2	34	4	0		38
	15.04	1.77	0.00		16.81
	89.47	10.53	0.00		
	18.78	20.00	0.00		
3	31	3	6		40
	13.72	1.33	2.65		17.70
	77.50	7.50	15.00		
	17.13	15.00	24.00		
4	23	3	7		33
	10.18	1.33	3.10		14.60
	69.70	9.09	21.21		
	12.71	15.00	28.00		
5	24	4	5		33
	10.62	1.77	2.21		14.60
	72.73	12.12	15.15		
	13.26	20.00	20.00		
6	39	3	1		43
	17.26	1.33	0.44		19.03
	90.70	6.98	2.33		
	21.55	15.00	4.00		
TOTAL	181	20	25		226
	80.09	8.85	11.06		100.00

CHI-SQUARE	14.569	DF=	10	PROB=0.1486
PHI	0.254			
CONTINGENCY COEFFICIENT	0.246			
CRAMER'S V	0.180			
LIKELIHOOD RATIO CHISQUARE	19.521	DF=	10	PROB=0.0364

ALL GROUPS

TABLE OF GROUP BY N9

GROUP	FINAL GROUP	N9	EXERCISE	
FREQUENCY				
PERCENT				
ROW PCT				
COL PCT	HIGH	MEDIUM	LOW	TOTAL
1	3	1	35	39
	1.33	0.44	15.49	17.26
	7.69	2.56	89.74	
	12.50	14.29	17.95	
2	4	1	33	38
	1.77	0.44	14.60	16.81
	10.53	2.63	86.84	
	16.67	14.29	16.92	
3	7	2	31	40
	3.10	0.88	13.72	17.70
	17.50	5.00	77.50	
	29.17	28.57	15.90	
4	4	2	27	33
	1.77	0.88	11.95	14.60
	12.12	6.06	81.82	
	16.67	28.57	13.85	
5	3	0	30	33
	1.33	0.00	13.27	14.60
	9.09	0.00	90.91	
	12.50	0.00	15.38	
6	3	1	39	43
	1.33	0.44	17.26	19.03
	6.98	2.33	90.70	
	12.50	14.29	20.00	
TOTAL	24	7	195	226
	10.62	3.10	86.28	100.00

CHI-SQUARE 6.017 DF= 10 PROB=0.6138
 PHI 0.163
 CONTINGENCY COEFFICIENT 0.161
 CRAMER'S V 0.115
 LIKELIHOOD RATIO CHISQUARE 6.595 DF= 10 PROB=0.7630

ALL GROUPS

TABLE OF GROUP BY N10

GROUP	FINAL GROUP	N10	HEART ATTACK	
FREQUENCY				
PERCENT				
ROW PCT				
COL PCT	HIGH	MEDIUM	LOW	TOTAL
1	3	1	35	39
	1.33	0.44	15.49	17.26
	7.69	2.56	89.74	
	18.75	16.67	17.16	
2	2	0	36	38
	0.88	0.00	15.93	16.81
	5.26	0.00	94.74	
	12.50	0.00	17.65	
3	3	1	36	40
	1.33	0.44	15.93	17.70
	7.50	2.50	90.00	
	18.75	16.67	17.65	
4	2	1	30	33
	0.88	0.44	13.27	14.60
	6.06	3.03	90.91	
	12.50	16.67	14.71	
5	3	1	29	33
	1.33	0.44	12.83	14.60
	9.09	3.03	87.88	
	18.75	16.67	14.22	
6	3	2	38	43
	1.33	0.88	16.81	19.03
	6.98	4.65	88.37	
	18.75	33.33	18.63	
TOTAL	16	6	204	226
	7.08	2.65	90.27	100.00

CHI-SQUARE 2.265 DF= 10 PROB=0.9939
 PHI 0.100
 CONTINGENCY COEFFICIENT 0.100
 CRAMER'S V 0.071
 LIKELIHOOD RATIO CHISQUARE 3.155 DF= 10 PROB=0.9775

ALL GROUPS

TABLE OF GROUP BY N11

GROUP	FINAL GROUP	N11	BELT/BP	
FREQUENCY				
PERCENT				
ROW PCT				
COL PCT	HIGH	MEDIUM	LOW	TOTAL
1	9	3	27	39
	3.98	1.33	11.95	17.26
	23.08	7.69	69.23	
	19.57	17.65	16.56	
2	11	3	24	38
	4.87	1.33	10.62	16.81
	28.95	7.89	63.16	
	23.91	17.65	14.72	
3	9	1	30	40
	3.98	0.44	13.27	17.70
	22.50	2.50	75.00	
	19.57	5.88	18.40	
4	9	4	20	33
	3.98	1.77	8.85	14.60
	27.27	12.12	60.61	
	19.57	23.53	12.27	
5	5	4	24	33
	2.21	1.77	10.62	14.60
	15.15	12.12	72.73	
	10.87	23.53	14.72	
6	3	2	38	43
	1.33	0.88	16.81	19.03
	6.98	4.65	88.37	
	6.52	11.76	23.31	
TOTAL	46	17	163	226
	20.35	7.52	72.12	100.00

CHI-SQUARE 12.981 DF= 10 PROB=0.2247
 PHI 0.240
 CONTINGENCY COEFFICIENT 0.233
 CRAMER'S V 0.169
 LIKELIHOOD RATIO CHISQUARE 14.281 DF= 10 PROB=0.1606

ALL GROUPS

TABLE OF GROUP BY N12

GROUP	FINAL GROUP N12			EYE CHECK
FREQUENCY PERCENT ROW PCT COL PCT	HIGH	MEDIUM	LOW	TOTAL
1	19 8.41 48.72 13.29	2 0.88 5.13 16.67	18 7.96 46.15 25.35	39 17.26
2	25 11.06 65.79 17.48	1 0.44 2.63 8.33	12 5.31 31.58 16.90	38 16.61
3	27 11.95 67.50 18.88	1 0.44 2.50 8.33	12 5.31 30.00 16.90	40 17.70
4	21 9.29 63.64 14.69	3 1.33 9.09 25.00	9 3.98 27.27 12.68	33 14.60
5	20 8.85 60.61 13.99	2 0.88 6.06 16.67	11 4.87 33.33 15.49	33 14.60
6	31 13.72 72.09 21.68	3 1.33 6.98 25.00	9 3.98 20.93 12.68	43 19.03
TOTAL	143 63.27	12 5.31	71 31.42	226 100.00

CHI-SQUARE	8.728	DF= 10	PROB=0.5561
PHI	0.197		
CONTINGENCY COEFFICIENT	0.193		
CRAMER'S V	0.139		
LIKELIHOOD RATIO CHISQUARE	8.768	DF= 10	PROB=0.5542

ALL GROUPS

TABLE OF GROUP BY N13

GROUP	FINAL	GROUP	N13	BELT/BP	
FREQUENCY					
PERCENT					
ROW PCT					
CCL PCT	HIGH	MEDIUM	LOW		TOTAL
1	7	1	31		39
	3.10	0.44	13.72		17.26
	17.95	2.56	79.49		
	21.86	12.50	16.67		
2	6	2	30		38
	2.65	0.88	13.27		16.81
	15.79	5.26	78.95		
	18.75	25.00	16.13		
3	3	2	35		40
	1.33	0.88	15.49		17.70
	7.50	5.00	87.50		
	9.38	25.00	18.82		
4	4	1	28		33
	1.77	0.44	12.39		14.60
	12.12	3.03	84.85		
	12.50	12.50	15.05		
5	5	1	27		33
	2.21	0.44	11.95		14.60
	15.15	3.03	81.82		
	15.63	12.50	14.52		
6	7	1	35		43
	3.10	0.44	15.49		19.03
	16.28	2.33	81.40		
	21.88	12.50	18.82		
TOTAL	32	8	186		226
	14.16	3.54	82.30		100.00

CHI-SQUARE 3.120 DF= 10 PROB=0.9785
 PHI 0.117
 CONTINGENCY COEFFICIENT 0.117
 CRAMER'S V 0.083
 LIKELIHOOD RATIO CHISQUARE 3.314 DF= 10 PROB=0.9731

ALL GROUPS

TABLE OF GROUP BY N14

GROUP	FINAL GROUP	N14	EATING	
FREQUENCY				
PERCENT				
ROW PCT				
COL PCT	HIGH	MEDIUM	LOW	TOTAL
1	14	1	24	39
	6.19	0.44	10.62	17.26
	35.90	2.56	61.54	
	16.67	5.88	19.20	
2	16	3	19	38
	7.08	1.33	8.41	16.81
	42.11	7.89	50.00	
	19.05	17.65	15.20	
3	14	7	19	40
	6.19	3.10	8.41	17.70
	35.00	17.50	47.50	
	16.67	41.18	15.20	
4	11	2	20	33
	4.87	0.88	8.85	14.60
	33.33	6.06	60.61	
	13.10	11.76	16.00	
5	13	2	18	33
	5.75	0.88	7.96	14.60
	39.39	6.06	54.55	
	15.48	11.76	14.40	
6	16	2	25	43
	7.08	0.88	11.06	19.03
	37.21	4.65	58.14	
	19.05	11.76	20.00	
TOTAL	84	17	125	226
	37.17	7.52	55.31	100.00

CHI-SQUARE	8.867	DF=	10	PROB=0.5448
PHI	0.198			
CONTINGENCY COEFFICIENT	0.194			
CRAMER'S V	0.140			
LIKELIHOOD RATIO CHISQUARE	7.948	DF=	10	PROB=0.6339

ALL GROUPS

TABLE OF GROUP BY N15

GROUP	FINAL GROUP	N15	DENTIST	
FREQUENCY				
PERCENT				
ROW PCT				
COL PCT	HIGH	MEDIUM	LOW	TOTAL
1	28	1	10	39
	12.39	0.44	4.42	17.26
	71.79	2.56	25.64	
	17.18	8.33	19.61	
2	28	1	9	38
	12.39	0.44	3.98	15.81
	73.68	2.63	23.68	
	17.18	8.33	17.65	
3	29	2	9	40
	12.83	0.88	3.98	17.70
	72.50	5.00	22.50	
	17.79	16.67	17.65	
4	26	3	4	33
	11.50	1.33	1.77	14.60
	78.79	9.09	12.12	
	15.95	25.00	7.64	
5	23	3	7	33
	10.18	1.33	3.10	14.60
	69.70	9.09	21.21	
	14.11	25.00	13.73	
6	29	2	12	43
	12.83	0.88	5.31	19.03
	67.44	4.65	27.91	
	17.79	16.67	23.53	
TOTAL	163	12	51	226
	72.12	5.31	22.57	100.00

CHI-SQUARE	5.612	DF=	10	PROB=0.8468
PHI	0.158			
CONTINGENCY COEFFICIENT	0.156			
CRAMER'S V	0.111			
LIKELIHOOD RATIO CHISQUARE	5.827	DF=	10	PROB=0.8296

ALL GROUPS

TABLE OF GROUP BY N16

GROUP	FINAL GROUP	N16	ACCIDENT/STR	
FREQUENCY				
PERCENT				
ROW PCT				
COL PCT	HIGH	MEDIUM	LOW	TOTAL
1	20	2	17	39
	8.85	0.88	7.52	17.26
	51.28	5.13	43.59	
	15.15	25.00	19.77	
2	26	1	11	38
	11.50	0.44	4.87	16.81
	68.42	2.63	28.95	
	19.70	12.50	12.79	
3	19	1	20	40
	8.41	0.44	8.85	17.70
	47.50	2.50	50.00	
	14.39	12.50	23.26	
4	19	2	12	33
	8.41	0.88	5.31	14.60
	57.58	6.06	36.36	
	14.39	25.00	13.95	
5	18	1	14	33
	7.96	0.44	6.19	14.60
	54.55	3.03	42.42	
	13.64	12.50	16.28	
6	30	1	12	43
	13.27	0.44	5.31	19.03
	69.77	2.33	27.91	
	22.73	12.50	13.95	
TOTAL	132	8	86	226
	58.41	3.54	38.05	100.00

CHI-SQUARE 8.125 DF= 10 PROB=0.6166
 PHI 0.190
 CONTINGENCY COEFFICIENT 0.186
 CRAMER'S V 0.134
 LIKELIHOOD RATIO CHISQUARE 8.656 DF= 10 PROB=0.6234

ALL GROUPS

TABLE OF GROUP BY N17

GROUP FINAL GROUP N17 LUNG CANCER

FREQUENCY PERCENT ROW PCT COL PCT	HIGH	MEDIUM	LOW	TOTAL
1	20 8.85 51.28 21.98	1 0.44 2.56 3.70	18 7.96 46.15 16.67	39 17.26
2	17 7.52 44.74 18.68	7 3.10 18.42 25.93	14 6.19 36.84 12.96	38 16.81
3	17 7.52 42.50 18.68	6 2.65 15.00 22.22	17 7.52 42.50 15.74	40 17.70
4	13 5.75 39.39 14.29	6 2.65 18.18 22.22	14 6.19 42.42 12.96	33 14.60
5	8 3.54 24.24 8.79	3 1.33 9.09 11.11	22 9.73 66.67 20.37	33 14.60
6	16 7.08 37.21 17.58	4 1.77 9.30 14.81	23 10.18 53.49 21.30	43 19.03
TOTAL	91 40.27	27 11.95	108 47.79	226 100.00

CHI-SQUARE 13.655 DF= 10 PROB=0.1797
 PHI 0.248
 CONTINGENCY COEFFICIENT 0.240
 CRAMER'S V 0.175
 LIKELIHOOD RATIO CHISQUARE 14.969 DF= 10 PROB=0.1332

ALL GROUPS

TABLE OF GROUP BY R

GROUP	FINAL GROUP			E			
FREQUENCY	PERCENT	ROW PCT	COL PCT	HIGH	MEDIUM	LOW	TOTAL
1	5	6	28	39			
	2.21	2.65	12.39	17.26			
	12.82	15.36	71.79				
	15.15	21.43	16.97				
2	6	4	28	38			
	2.65	1.77	12.39	16.81			
	15.79	10.53	73.68				
	18.18	14.29	16.97				
3	6	5	29	40			
	2.65	2.21	12.83	17.70			
	15.00	12.50	72.50				
	18.18	17.66	17.58				
4	7	1	25	33			
	3.10	0.44	11.06	14.60			
	21.21	3.03	75.76				
	21.21	3.57	15.15				
5	4	4	25	33			
	1.77	1.77	11.06	14.60			
	12.12	12.12	75.76				
	12.12	14.29	15.15				
6	5	8	30	43			
	2.21	3.54	13.27	19.03			
	11.63	18.60	69.77				
	15.15	28.57	18.18				
TOTAL	33	28	165	226			
	14.60	12.39	73.01	100.00			

CHI-SQUARE 5.720 DF= 10 PROB=0.8362
 PHI 0.159
 CONTINGENCY COEFFICIENT 0.157
 CRAMER'S V 0.112
 LIKELIHOOD RATIO CHISQUARE 6.461 DF= 10 PROB=0.7752

ALL GROUPS

TABLE OF R BY RACE

R	RACE		
FREQUENCY			
PERCENT			
ROW PCT			
COL PCT	A	W	TOTAL
HIGH	3	30	33
	1.33	12.27	14.60
	9.09	90.91	
	12.00	14.93	
MEDIUM	3	25	28
	1.33	11.06	12.39
	10.71	89.29	
	12.00	12.44	
LOW	19	146	165
	8.41	64.60	73.01
	11.52	80.48	
	76.00	72.64	
TOTAL	25	201	226
	11.06	86.94	100.00

CHI-SQUARE	0.168	DF=	2	PROB=0.9193
PHI	0.027			
CONTINGENCY COEFFICIENT	0.027			
CRAMER'S V	0.027			
LIKELIHOOD RATIO CHISQUARE	0.175	DF=	2	PROB=0.9161

ALL GROUPS
TABLE OF R BY SEX

R		SEX		
FREQUENCY				
PERCENT				
ROW PCT				
COL PCT	F	M		TOTAL
HIGH	28	5		33
	12.39	2.21		14.60
	84.85	15.15		
	16.28	9.26		
MEDIUM	19	9		28
	8.41	3.98		12.39
	67.86	32.14		
	11.05	16.67		
LOW	125	40		165
	55.31	17.70		73.01
	75.76	24.24		
	72.67	74.07		
TOTAL	172	54		226
	76.11	23.89		100.00

CHI-SQUARE	2.446	DF=	2	PROB=0.2944
PHI	0.104			
CONTINGENCY COEFFICIENT	0.103			
CRAMER'S V	0.104			
LIKELIHOOD RATIO CHISQUARE	2.524	DF=	2	PROB=0.2831

ALL GROUPS

TABLE OF R BY AGE

P		AGE					
FREQUENCY	PERCENT	ROW PCT	COL PCT	18-25	26-35	36+	TOTAL
-----+-----+-----+-----+-----							
HIGH				12	10	11	33
				5.31	4.42	4.87	14.60
				36.36	30.30	33.33	
				17.91	11.63	15.07	
-----+-----+-----+-----+-----							
MEDIUM				9	7	12	28
				3.98	3.10	5.31	12.39
				32.14	25.00	42.86	
				13.43	8.14	16.44	
-----+-----+-----+-----+-----							
LOW				46	69	50	165
				20.35	30.53	22.12	73.01
				27.88	41.82	30.30	
				68.66	80.23	68.49	
-----+-----+-----+-----+-----							
TOTAL				67	86	73	226
				29.65	38.05	32.30	100.00

CHI-SQUARE	4.505	DF=	4	PROB=0.3663
PHI	0.138			
CONTINGENCY COEFFICIENT	0.137			
CRAMER'S V	0.098			
LIKELIHOOD RATIO CHISQUARE	4.581	DF=	4	PROB=0.3569

GROUPS 1-5

TABLE OF N5 BY RACE

N5	BELT/EP		RACE
FREQUENCY			
PERCENT			
ROW PCT			
COL PCT	A	W	TOTAL
HIGH	13	33	46
	7.10	18.03	25.14
	28.26	71.74	
	61.90	20.37	
MEDIUM	1	11	12
	0.55	6.01	6.56
	8.33	91.67	
	4.76	6.79	
LOW	7	118	125
	3.83	64.48	68.31
	5.60	94.40	
	33.33	72.84	
TOTAL	21	162	183
	11.48	88.52	100.00

CHI-SQUARE	17.123	DF=	2	PROB=0.0002
PHI	0.306			
CONTINGENCY COEFFICIENT	0.293			
CRAMER'S V	0.306			
LIKELIHOOD RATIO CHISQUARE	14.806	DF=	2	PROB=0.0006

GROUPS 1-5

TABLE OF N5 BY SEX

N5	DELTA/PP		SE
FREQUENCY			
PERCENT			
ROW PCT			
COL PCT	F	M	TOTAL
HIGH	36	8	46
	20.77	4.37	25.14
	82.61	17.39	
	27.14	18.60	
MEDIUM	8	4	12
	4.37	2.19	6.56
	66.67	33.33	
	5.71	9.30	
LOW	94	31	125
	51.37	16.94	68.31
	75.20	24.80	
	67.14	72.09	
TOTAL	140	43	183
	76.50	23.50	100.00

CHI-SQUARE	1.718	DF=	2	PROB=0.4236
PHI	0.097			
CONTINGENCY COEFFICIENT	0.096			
CRAMER'S V	0.097			
LIKELIHOOD RATIO CHISQUARE	1.733	DF=	2	PROB=0.4204

GROUPS 1-5

TABLE OF N5 BY AGE

N5	BELT/BP		AGE		
FREQUENCY	PERCENT		ROW PCT		
COL PCT	16-25	26-35	36+	TOTAL	
HIGH	14	14	18	46	
	7.65	7.65	9.84	25.14	
	30.43	30.43	39.13		
	28.00	19.44	29.51		
MEDIUM	4	5	3	12	
	2.19	2.73	1.64	6.56	
	33.33	41.67	25.00		
	8.00	6.94	4.92		
LOW	32	53	40	125	
	17.49	28.96	21.86	68.31	
	25.60	42.40	32.00		
	64.00	73.61	65.57		
TOTAL	50	72	61	183	
	27.32	39.34	33.33	100.00	

CHI-SQUARE	2.479	DF=	4	PROB=0.6484
PHI	0.116			
CONTINGENCY COEFFICIENT	0.116			
CRAMER'S V	0.082			
LIKELIHOOD RATIO CHISQUARE	2.537	DF=	4	PROB=0.6380

GROUPS 1-5

TABLE OF N5 BY R

N5	BELT/RP			R	
FREQUENCY					
PERCENT					
ROW PCT					
COL PCT	HIGH	MEDIUM	LOW		TOTAL
HIGH	9	5	32		46
	4.92	2.73	17.49		25.14
	19.57	10.87	69.57		
	32.14	25.00	23.70		
MEDIUM	1	0	11		12
	0.55	0.00	6.01		6.56
	8.33	0.00	91.67		
	3.57	0.00	8.15		
LOW	18	15	92		125
	9.84	8.20	50.27		68.31
	14.40	12.00	73.60		
	64.29	75.00	68.15		
TOTAL	28	20	135		183
	15.30	10.93	73.77		100.00

CHI-SQUARE	3.068	DF=	4	PROB=0.5465
PHI	0.129			
CONTINGENCY COEFFICIENT	0.128			
CRAMER'S V	0.092			
LIKELIHOOD RATIO CHISQUARE	4.373	DF=	4	PROB=0.3579

GROUPS 1-5

TABLE OF N5 BY N2

N5	BELT/BP	N2	CHECKUP	
FREQUENCY				
PERCENT				
ROW PCT				
COL PCT	HIGH	MEDIUM	LOW	TOTAL
HIGH	13	7	26	46
	7.10	3.83	14.21	25.14
	28.26	15.22	56.52	
	35.14	58.33	19.40	
MEDIUM	3	1	8	12
	1.64	0.55	4.37	6.56
	25.00	8.33	66.67	
	8.11	8.33	5.97	
LOW	21	4	100	125
	11.48	2.19	54.64	68.31
	16.80	3.20	80.00	
	56.76	33.33	74.63	
TOTAL	37	12	134	183
	20.22	6.56	73.22	100.00

CHI-SQUARE	12.404	DF=	4	PROB=0.0146
PHI	0.260			
CONTINGENCY COEFFICIENT	0.252			
CRAMER'S V	0.184			
LIKELIHOOD RATIO CHISQUARE	11.463	DF=	4	PROB=0.0218

GROUPS 1-5

TABLE OF N5 BY N6

N5	BELT/BP	N6	SMOKING	
FREQUENCY				
PERCENT				
ROW PCT				
COL PCT	HIGH	MEDIUM	LOW	TOTAL
HIGH	3	2	41	46
	1.64	1.09	22.40	25.14
	6.52	4.35	89.13	
	21.43	33.33	25.15	
MEDIUM	0	1	11	12
	0.00	0.55	6.01	6.56
	0.00	8.33	91.67	
	0.00	16.67	6.75	
LOW	11	3	111	125
	6.01	1.64	60.66	68.31
	8.80	2.40	88.80	
	78.57	50.00	68.10	
TOTAL	14	6	163	183
	7.65	3.28	89.07	100.00

CHI-SQUARE	2.611	DF=	4	PROB=0.6250
PHI	0.119			
CONTINGENCY COEFFICIENT	0.119			
CRAMER'S V	0.084			
LIKELIHOOD RATIO CHISQUARE	3.255	DF=	4	PROB=0.5160

GROUPS 1-5

TABLE OF N11 BY RACE

N11	BELT/PP		RACE
FREQUENCY			
PERCENT			
ROW PCT			
COL PCT	A	W	TOTAL
HIGH	4	39	43
	2.19	21.31	23.50
	9.30	90.70	
	19.05	24.07	
MEDIUM	3	12	15
	1.64	6.56	8.20
	20.00	80.00	
	14.29	7.41	
LOW	14	111	125
	7.65	60.66	68.31
	11.20	88.80	
	66.67	68.52	
TOTAL	21	162	183
	11.48	88.52	100.00

CHI-SQUARE	1.282	DF=	2	PROB=0.5267
PHI	0.084			
CONTINGENCY COEFFICIENT	0.083			
CFAMER'S V	0.084			
LIKELIHOOD RATIO CHISQUARE	1.125	DF=	2	PROB=0.5699

GROUPS 1-5

TABLE OF N11 BY SEX

N11	BELT/PP		SEA	
FREQUENCY				
PERCENT				
ROW PCT				
COL PCT	F	M		TOTAL
HIGH	33	10		43
	18.03	5.46		23.50
	76.74	23.26		
	23.57	23.26		
MEDIUM	10	5		15
	5.46	2.73		8.20
	66.67	33.33		
	7.14	11.63		
LOW	97	28		125
	53.01	15.30		68.31
	77.60	22.40		
	69.29	65.12		
TOTAL	140	43		183
	76.50	23.50		100.00

CHI-SQUARE	0.892	DF=	2	PROB=0.6400
PHI	0.070			
CONTINGENCY COEFFICIENT	0.070			
CRAMER'S V	0.070			
LIKELIHOOD RATIO CHISQUARE	0.830	DF=	2	PROB=0.6602

GROUPS 1-5

TABLE OF N11 BY AGE

N11	BELT/EP			AGE	
FREQUENCY					
PERCENT					
ROW PCT					
COL PCT	18-25	26-35	36+		TOTAL
HIGH	17	16	10		43
	9.29	8.74	5.46		23.50
	39.53	37.21	23.26		
	34.00	22.22	16.39		
MEDIUM	2	6	7		15
	1.09	3.28	3.83		8.20
	13.33	40.00	46.67		
	4.00	8.33	11.48		
LOW	31	50	44		125
	16.94	27.32	24.04		68.31
	24.80	40.00	35.20		
	62.00	69.44	72.13		
TOTAL	50	72	61		183
	27.32	39.34	33.33		100.00

CHI-SQUARE	6.019	DF=	4	PROB=0.1978
PHI	0.181			
CONTINGENCY COEFFICIENT	0.178			
CRAMER'S V	0.128			
LIKELIHOOD RATIO CHISQUARE	6.058	DF=	4	PROB=0.1948

GROUPS 1-5

TABLE OF N11 BY F

N11	BELT/BP			F
FREQUENCY				
PERCENT				
ROW PCT				
COL PCT	HIGH	MEDIUM	LOW	TOTAL
HIGH	8	2	33	43
	4.37	1.09	18.03	23.50
	18.60	4.65	76.74	
	28.57	10.00	24.44	
MEDIUM	2	3	10	15
	1.09	1.64	5.46	8.20
	13.33	20.00	66.67	
	7.14	15.00	7.41	
LOW	18	15	92	125
	9.84	8.20	50.27	68.31
	14.40	12.00	73.60	
	64.29	75.00	68.15	
TOTAL	28	20	135	183
	15.30	10.93	73.77	100.00

CHI-SQUARE	3.577	DF=	4	PROB=0.4969
PHI	0.136			
CONTINGENCY COEFFICIENT	0.135			
CRAMER'S V	0.096			
LIKELIHOOD RATIO CHISQUARE	3.565	DF=	4	PROB=0.4681

GROUPS 1-5

TABLE OF N11 BY N2

N11	BELT/BP	N2	CHECKUP	
FREQUENCY				
PERCENT				
ROW PCT				
COL PCT	HIGH	MEDIUM	LOW	TOTAL
HIGH	10	4	29	43
	5.46	2.19	15.55	23.50
	23.26	9.30	67.44	
	27.03	33.33	21.64	
MEDIUM	4	2	9	15
	2.19	1.09	4.92	8.20
	26.67	13.33	60.00	
	10.81	16.67	6.72	
LOW	23	6	96	125
	12.57	3.28	52.46	68.31
	18.40	4.80	76.80	
	62.16	50.00	71.64	
TOTAL	37	12	134	183
	20.22	6.56	73.22	100.00

CHI-SQUARE	3.615	DF=	4	PROB=0.4606
PHI	0.141			
CONTINGENCY COEFFICIENT	0.139			
CRAMER'S V	0.099			
LIKELIHOOD RATIO CHISQUARE	3.369	DF=	4	PROB=0.4961

GROUPS 1-5

TABLE OF N11 BY N6

N11	BELT/BP	N6	SMCKING	
FREQUENCY				
PERCENT				
ROW PCT				
COL PCT	HIGH	MEDIUM	LOW	TOTAL
HIGH	1	3	39	43
	0.55	1.64	21.31	23.50
	2.33	6.98	90.70	
	7.14	50.00	23.93	
MEDIUM	4	1	10	15
	2.19	0.55	5.46	8.20
	26.67	6.67	66.67	
	28.57	16.67	6.13	
LOW	9	2	114	125
	4.92	1.09	62.30	68.31
	7.20	1.60	91.20	
	64.29	33.33	69.94	
TOTAL	14	6	163	183
	7.65	3.28	89.07	100.00

CHI-SQUARE	13.032	DF=	4	PROB=0.0111
PHI	0.267			
CONTINGENCY COEFFICIENT	0.258			
CRAMER'S V	0.189			
LIKELIHOOD RATIO CHISQUARE	10.599	DF=	4	PROB=0.0315

GROUPS 1-5

TABLE OF N16 BY RACE

N16	ACCIDENT/STR	RACE	
FREQUENCY			
PERCENT			
ROW PCT			
COL PCT	N	W	TOTAL
HIGH	7	95	102
	3.83	51.91	55.74
	6.86	93.14	
	33.33	56.64	
MEDIUM	0	7	7
	0.00	3.83	3.83
	0.00	100.00	
	0.00	4.32	
LOW	14	60	74
	7.65	32.79	40.44
	18.92	81.08	
	66.67	37.04	
TOTAL	21	162	183
	11.46	86.52	100.00

CHI-SQUARE	7.680	DF=	2	PROB=0.0290
PHI	0.197			
CONTINGENCY COEFFICIENT	0.193			
CRAMER'S V	0.197			
LIKELIHOOD RATIO CHISQUARE	7.619	DF=	2	PROB=0.0222

GROUPS 1-5

TABLE OF N16 BY SEX

N16	ACCIDENT/STRE		SEX
FREQUENCY			
PFRCENT			
ROW PCT			
COL PCT	F	M	TOTAL
HIGH	81	21	102
	44.26	11.48	55.74
	79.41	20.59	
	57.86	46.84	
MEDIUM	2	5	7
	1.09	2.73	3.83
	26.57	71.43	
	1.43	11.63	
LOW	57	17	74
	31.15	9.29	40.44
	77.03	22.97	
	40.71	39.53	
TOTAL	140	43	183
	76.50	23.50	100.00

CHI-SQUARE	9.438	DF=	2	PROB=0.0089
PHI	0.227			
CONTINGENCY COEFFICIENT	0.221			
CRAMER'S V	0.227			
LIKELIHOOD RATIO CHISQUARE	7.685	DF=	2	PROB=0.0214

GROUPS 1-5

TABLE OF N16 BY AGE

N16	ACCIDENT/STR			AGE	
FREQUENCY					
PERCENT					
ROW PCT					
COL PCT	18-25	26-35	36+		TOTAL
HIGH	24	41	37		102
	13.11	22.40	20.22		55.74
	23.53	40.20	36.27		
	48.00	56.94	60.66		
MEDIUM	0	5	2		7
	0.00	2.73	1.09		3.83
	0.00	71.43	28.57		
	0.00	6.94	3.28		
LOW	26	26	22		74
	14.21	14.21	12.02		40.44
	35.14	35.14	29.73		
	52.00	36.11	36.07		
TOTAL	50	72	61		183
	27.32	39.34	33.33		100.00

CHI-SQUARE	6.687	DF=	4	PROB=0.1420
PHI	0.194			
CONTINGENCY COEFFICIENT	0.190			
CRAMER'S V	0.137			
LIKELIHOOD RATIO CHISQUARE	8.347	DF=	4	PROB=0.0797

GROUPS 1-5

TABLE OF N16 BY R

N16	ACCIDENT/STR			R
FREQUENCY				
PERCENT				
ROW PCT				
COL PCT	HIGH	MEDIUM	LOW	TOTAL
HIGH	11	13	78	102
	6.01	7.10	42.62	55.74
	10.78	12.75	76.47	
	39.29	65.00	57.78	
MEDIUM	1	0	6	7
	0.55	0.00	3.28	3.83
	14.29	0.00	85.71	
	3.57	0.00	4.44	
LOW	16	7	51	74
	8.74	3.63	27.87	40.44
	21.62	9.46	68.92	
	57.14	35.00	37.78	
TOTAL	28	20	135	183
	15.30	10.93	73.77	100.00

CHI-SQUARE	4.988	DF=	4	PROB=0.2885
PHI	0.165			
CONTINGENCY COEFFICIENT	0.163			
CRAMER'S V	0.117			
LIKELIHOOD RATIO CHISQUARE	5.678	DF=	4	PROB=0.2245

GROUPS 1-5

TABLE OF N16 BY N2

N16	ACCIDENT/STR			N2	CHECKUP
FREQUENCY PERCENT ROW PCT COL PCT	HIGH	MEDIUM	LOW	TOTAL	
HIGH	10 5.46 9.80 27.03	4 2.19 3.92 33.33	88 48.09 86.27 65.67	102 55.74	
MEDIUM	3 1.64 42.86 8.11	0 0.00 0.00 0.00	4 2.19 57.14 2.99	7 3.83	
LOW	24 13.11 32.43 64.86	8 4.37 10.81 66.67	42 22.95 56.76 31.34	74 40.44	
TOTAL	37 20.22	12 6.56	134 73.22	183 100.00	

CHI-SQUARE	21.648	DF=	4	PROB=0.0002
PHI	0.344			
CONTINGENCY COEFFICIENT	0.325			
CRAMER'S V	0.243			
LIKELIHOOD RATIO CHISQUARE	22.084	DF=	4	PROB=0.0002

GROUPS 1-5

TABLE OF N16 BY N6

N16	ACCIDENT/STR	No	SMCKING	
FREQUENCY	PERCENT	ROW PCT	COL PCT	TOTAL
	HIGH	MEDIUM	LOW	
HIGH	7	3	92	102
	3.83	1.64	50.27	55.74
	6.86	2.94	90.20	
	50.00	50.00	56.44	
MEDIUM	0	0	7	7
	0.00	0.00	3.83	3.83
	0.00	0.00	100.00	
	0.00	0.00	4.29	
LOW	7	3	64	74
	3.63	1.64	34.97	40.44
	9.46	4.05	86.49	
	50.00	50.00	39.26	
TOTAL	14	6	163	183
	7.65	3.28	89.07	100.00

CHI-SQUARE 1.499 DF= 4 PROB=0.8268
 PHI 0.091
 CONTINGENCY COEFFICIENT 0.090
 CRAMER'S V 0.064
 LIKELIHOOD RATIO CHISQUARE 2.234 DF= 4 PROB=0.6929

REFERENCES

- Ashton, S. J., Mackay, G. M., and Camm, S. Seat belt use in Britain under voluntary and mandatory conditions. 27th Proceedings American Association for Automotive Medicine, Arlington Heights, IL, 1983.
- Schwalm, Norman D. and Slovic, Paul. Risk Perception and Seat Belt Use: Report on Phase II: Experimental Test of Animatic Equivalent Seat Belt Messages, US DOT, NHTSA, 1983.
- Slovic, Paul, Lichtenstein, Sarah, MacGregor, Don, Fischhoff, Baruch, and Schwalm, Norman. Risk Perception and Seat-Belt Use: Report on Phase I: Development and Test of Printed Seat-Belt Messages, US DOT, NHTSA, 1983.